

# Vectors

## Introduction:

In our real life situation we deal with physical quantities such as distance, speed, temperature, volume etc. These quantities are sufficient to describe change of position, rate of change of position, body temperature or temperature of a certain place and space occupied in a confined portion respectively.

We have also come across physical quantities such as displacement, velocity, acceleration, momentum etc, which are of different type in comparison to above.

Consider the figure-1, where A, B, C are

at a distance 4 km from P. If we start from P, then covering 4 km distance is not sufficient to

describe the destination where we reach after the travel, so

here the end point plays an important role giving rise the need of direction. So we need to

study about direction of a quantity, along with magnitude.




Fig - 1

## Objective

After completion of the topic you are able to:-

- (i) Define and distinguish between scalars and vectors.
- (ii) Represent a vector as directed line segment.
- (iii) classify vectors into different types.
- (iv) Resolve vector along two or three mutually perpendicular axes.
- (v) Define dot product of two vectors and explain its geometrical meaning.
- (vi) Define cross product of two vectors and apply it to find area of triangle and parallelogram.

# Scalars and Vectors

All the physical quantities can be divided into two types

- (i) Scalar quantity or Scalars
- (ii) Vector quantity or Vectors

Scalar quantity:- The physical quantities which require only magnitude for its complete specification is called as Scalar quantities.

Examples:- Speed, mass, distance, velocity, volume etc.

Vector:- A directed line segment is called as Vector.

Vector quantities:- A physical quantity which requires both magnitude & direction for its complete specification and satisfies the law of vector addition is called as Vector quantities.

Examples:- Displacement, Force, acceleration, velocity, momentum etc.

Representation of Vector:- A vector is directed line segment  $\overrightarrow{AB}$  where A is the initial point and B is the terminal point and direction is from A to B. (See Fig-2)

Similarly  $\overrightarrow{BA}$  is a directed line which represents a vector having initial point B and Terminal A.




Fig-2



Fig-3

Notation:- A vector quantity is always represented by an arrow ( $\rightarrow$ ) mark over it or by bars ( $-$ ) over it. For example  $\overrightarrow{AB}$ . It is also represented by a single small letter with an arrow or bars mark over it. For example  $\vec{a}$ .

Magnitude of  $\overrightarrow{AB} = |\overrightarrow{AB}| = \text{Length } AB = AB$ .



Types of Vector :- Vectors are following types

- (1) Null Vector or Zero Vector or Void Vector - A vector having zero magnitude and arbitrary direction is called as a null vector and is denoted by  $\vec{0}$ .

Clearly, a null vector has no definite direction. If  $\vec{a} = \vec{AB}$ , then  $\vec{a}$  is a null (or zero) vector if  $|\vec{a}| = 0$  i.e. if  $|\vec{AB}| = 0$

- (2) Proper Vector - Any non zero vector is called as a proper vector. If  $|\vec{a}| \neq 0$  then  $\vec{a}$  is a proper vector.


- (3) Unit Vector - A vector whose magnitude is unity is called a unit vector. Unit vectors are denoted by a small letter <sup>1</sup> over it. For example  $\hat{a}$ .  $|\hat{a}| = 1$ .

**Note** :- The unit vector along the direction of a vector  $\vec{a}$  is given by

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

- (4) Co-initial vectors :- Vectors having the same initial point are called co-initial vector.

In figure-4,  $\vec{OA}$ ,  $\vec{OB}$ ,  $\vec{OC}$ ,  $\vec{OD}$  and  $\vec{OE}$  are co-initial vectors.



- (5) Like and Unlike Vectors :- Vectors are said to be like if they have same direction and unlike if they have opposite direction.

6) Co-Linear Vectors:- Vectors are said to be co-linear or parallel if they have the same line of action. In figure-5  $\vec{AB}$  and  $\vec{BC}$  are co-linear.




Fig 5

7) Parallel Vectors:- Vectors are said to be parallel if they have same line of action or have line of action parallel to one another. In fig-6 the vectors are parallel to each other.




Fig 6

8) Co-planer Vectors:- Vectors are said to be co-planer if they lie on the same plane. In fig-7 vector  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are coplaner.




Fig-7

9) Negative of a Vector:- A vector having same magnitude but opposite in direction to that of a given vector is called negative vector of that vector. If  $\vec{a}$  is any vector then negative vector of it is written as  $-\vec{a}$  and  $|\vec{a}| = |-\vec{a}|$  but both have direction opposite to each other as shown in fig-8.





Fig-8.

10) Equal Vectors:- Two vectors are said to be equal if they have same magnitude as well as same direction.



Thus  $\vec{a} = \vec{b}$ .

Notes :- Two vectors can not be equal

- (i) If they have different magnitude
- (ii) If they have inclined supports
- (iii) If they have different Sense.



# Vector operations

## Addition of vectors:-

Triangle law of vector addition:- The law states that if two vectors are represented by the two sides of a triangle taken in same order their sum or resultant is represented by the 3<sup>rd</sup> side of the triangle with direction in reverse order.

As shown in Figure-10  $\vec{a}$  and  $\vec{b}$  are two vectors represented by two sides OA and AB of a triangle ABC in same order. Then the sum  $\vec{a} + \vec{b}$  is represented by the third side OB taken in reverse order. i.e. the vector  $\vec{a}$  is represented by the directed segment  $\vec{OA}$  and the vector  $\vec{b}$  be the directed segment  $\vec{AB}$ , so that the terminal point A of  $\vec{a}$  is the initial point of  $\vec{b}$ . Then  $\vec{OB}$  represents the sum (or resultant)  $(\vec{a} + \vec{b})$ . Thus  $\vec{OB} = \vec{a} + \vec{b}$




Fig-10

Note-1- The method of drawing of triangle in order to define the vector sum  $(\vec{a} + \vec{b})$  is called triangle law of addition of the vectors.

Note-2- Since any side of a triangle is less than the sum of the other two sides.  $|\vec{OB}| \neq |\vec{OA}| + |\vec{AB}|$

Parallelogram law of vector addition-

If  $\vec{a}$  and  $\vec{b}$  are two vectors represented by two adjacent sides of a parallelogram in a magnitude and direction, then their sum (resultant) is represented in magnitude and direction by the diagonal which is passing through the common initial point of the two vectors. As shown in fig-11 if OA is  $\vec{a}$  and AB is  $\vec{b}$  then OB diagonal represent  $\vec{a} + \vec{b}$ .




Fig-11

$$\text{i.e. } \vec{a} + \vec{b} = \vec{OA} + \vec{AB}$$




Fig-12

Polygon law of vector addition:- If  $\vec{a}, \vec{b}, \vec{c}$  and  $\vec{d}$  are the four sides of a polygon in the same order then their sum is represented by the last side of the polygon taken in opposite order as shown in Fig-12.

### Subtraction of two Vectors

If  $\vec{a}$  and  $\vec{b}$  are two given vectors then the subtraction of  $\vec{b}$  from  $\vec{a}$  denoted by  $\vec{a} - \vec{b}$  is defined as addition of  $-\vec{b}$  with  $\vec{a}$ . i.e.,  $\vec{a} - \vec{b} = \vec{a} + (-\vec{b})$

Properties of vector addition:-

- (i) Vector addition is commutative i.e., if  $\vec{a}$  &  $\vec{b}$  are any two vectors then  $\vec{a} + \vec{b} = \vec{b} + \vec{a}$
- (ii) Vector addition is associative i.e., if  $\vec{a}, \vec{b}, \vec{c}$  are any three vectors then  $(\vec{a} + \vec{b}) + \vec{c} = \vec{a} + (\vec{b} + \vec{c})$
- (iii) Existence of additive identity i.e., for any vector  $\vec{a}$ ,  $\vec{0}$  is the additive identity i.e.,  $\vec{0} + \vec{a} = \vec{a} + \vec{0} = \vec{a}$  where  $\vec{0}$  is a null vector.
- (iv) Existence of additive inverse:- If  $\vec{a}$  is any non-zero vector then  $-\vec{a}$  is the additive inverse of  $\vec{a}$  so that  $\vec{a} + (-\vec{a}) = (-\vec{a}) + \vec{a} = \vec{0}$



## Multiplication of a vector by a scalar

If  $\vec{a}$  is a vector and  $K$  is a non-zero scalar then the multiplication of the vector  $\vec{a}$  by the scalar  $K$  is a vector denoted by  $K\vec{a}$  or  $\vec{a}K$  whose magnitude is  $|K|$  times that of  $\vec{a}$ .

$$\text{i.e. } K\vec{a} = |K| \times |\vec{a}|$$

$$= K \times |\vec{a}| \text{ if } K \geq 0$$

$$= (-K) \times |\vec{a}| \text{ if } K < 0$$

The direction of  $K\vec{a}$  is same as that of  $\vec{a}$  if  $K$  is positive and opposite as that of  $\vec{a}$  if  $K$  is negative.  $K\vec{a}$  and  $\vec{a}$  are always parallel to each other.

### Properties of scalar multiplication of Vectors:-

If  $h$  and  $K$  are scalars and  $\vec{a}$  and  $\vec{b}$  are given vectors then

- (i)  $K(\vec{a} + \vec{b}) = K\vec{a} + K\vec{b}$
- (ii)  $(h + K)\vec{a} = h\vec{a} + K\vec{a}$ , (Distributive law)
- (iii)  $(hK)\vec{a} = h(K\vec{a})$ , (Associative law)
- (iv)  $1 \cdot \vec{a} = \vec{a}$
- (v)  $0 \cdot \vec{a} = \vec{0}$ .

## Position vector of a point

Let  $O$  be a fixed point called origin, let  $P$  be any other point, then the vector  $\vec{OP}$  is called position vector of the point  $P$  relative to  $O$  and is denoted by  $\vec{P}$ .

As shown in figure - 13, let  $AB$  be any vector, then applying

triangle law of addition we have  $\vec{OA} + \vec{AB} = \vec{OB}$  where  $\vec{OA}$

$$= \vec{a} \text{ and } \vec{OB} = \vec{b} \Rightarrow \vec{AB} = \vec{OB} - \vec{OA} = \vec{b} - \vec{a}$$

$$= (\text{position vector of } B) - (\text{position vector of } A)$$

Section Formula :- Let  $A$  and  $B$  be two points with position vector  $\vec{a}$  and  $\vec{b}$  respectively

and  $P$  be a point on line segment  $AB$ , dividing it in

the ratio  $m:n$  internally. Then the position vector of  $P$  i.e.  $\vec{r}$  is given by the formula:

$$\vec{r} = \frac{m\vec{b} + n\vec{a}}{m+n}$$

If  $P$  divides  $AB$  externally in the ratio  $m:n$  then

$$\vec{r} = \frac{m\vec{b} - n\vec{a}}{m-n}$$

If  $P$  is the midpoint of  $AB$  then  $\vec{r} = \frac{\vec{a} + \vec{b}}{2}$

Example - 1 :- prove that by vector method the medians of a triangle are concurrent.

Solution :- Let  $ABC$  be a triangle where  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are the position vector of  $A$ ,  $B$  and  $C$  respectively. We have to show that the medians of this triangle are concurrent.




Fig - 13




Fig - 14




Fig - 15



Let AD, BE and CF are the three medians of the triangle. Now as D be the midpoint of BC, So position vector of D i.e.  $\vec{d} = \frac{\vec{b} + \vec{c}}{2}$ .

Let G be any point of the median AD which divides AD in the ratio 2:1. Then position vector of G is given by  $\vec{g} = \frac{2\vec{d} + \vec{a}}{2+1} = \frac{2(\frac{\vec{b} + \vec{c}}{2}) + 1\vec{a}}{3}$

(by applying section formula)

$$\Rightarrow \vec{g} = \frac{\vec{a} + \vec{b} + \vec{c}}{3}$$



Let  $G'$  be point which divides  $BE$  in the ratio  $2:1$

Position vector of  $E$  is  $\vec{e} = \frac{\vec{a} + \vec{c}}{2}$ .

Then position vector of  $G'$  is given by  $\vec{g}' = \frac{2\vec{e} + \vec{b}}{2+1} = \frac{2(\frac{\vec{a} + \vec{c}}{2}) + \vec{b}}{3}$

$$\Rightarrow \vec{g}' = \frac{\vec{a} + \vec{b} + \vec{c}}{3} \quad \text{(by applying section formula)}$$

As position vector of a point is unique, so  $G = G'$   
 Similarly if we take  $G''$  be a point on  $CF$  dividing it in  $2:1$  ratio then the position vector of  $G''$  will be same as that of  $G$ .

Hence  $G$  is the one point where three medians meet.

$\therefore$  The three medians of a triangle are concurrent. (Proved)

Example 2: Prove that (i)  $|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$  (it is known as triangle inequality).

$$(ii) |\vec{a}| - |\vec{b}| \leq |\vec{a} - \vec{b}|$$

$$(iii) |\vec{a} - \vec{b}| \leq |\vec{a}| + |\vec{b}|$$

Proof: - Let  $O, A$  and  $B$  be three points, which are not collinear and then draw a triangle

$OAB$ .

Let  $\vec{OA} = \vec{a}$ ,  $\vec{AB} = \vec{b}$ , then by triangle law of addition we have  $\vec{OB} = \vec{a} + \vec{b}$

From properties of triangle we know that the sum of any two sides of a triangle is greater than the third side.




Fig - 16

$$\Rightarrow OB < OA + AB$$

$$\Rightarrow |\vec{OB}| < |\vec{OA}| + |\vec{AB}|$$

$$\Rightarrow |\vec{a} + \vec{b}| < |\vec{a}| + |\vec{b}| \quad \text{--- (1)}$$

When  $O, A, B$  are collinear then from Fig-17 it is clear that  $OB = OA + AB$

$$\Rightarrow |\vec{OB}| = |\vec{OA}| + |\vec{AB}|$$

$$\Rightarrow |\vec{a} + \vec{b}| = |\vec{a}| + |\vec{b}| \quad \text{--- (2)}$$



Fig-17



From (1) and (2) we have

$$|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}| \text{ (proved)}$$

$$(i) |\vec{a}| = |\vec{a} - \vec{b} + \vec{b}| \text{ ----- (1)}$$

But  $|\vec{a} - \vec{b} + \vec{b}| \leq |\vec{a} - \vec{b}| + |\vec{b}|$  (from triangle inequality) - (2)

From (1) and (2) we get  $|\vec{a}| \leq |\vec{a} - \vec{b}| + |\vec{b}|$

$$\Rightarrow |\vec{a}| - |\vec{b}| \leq |\vec{a} - \vec{b}| \text{ (proved)}$$

$$(ii) |\vec{a} - \vec{b}| = |\vec{a} + (-\vec{b})| \leq |\vec{a}| + |-\vec{b}| \text{ (from triangle inequality)} \\ = |\vec{a}| + |\vec{b}| \text{ (as } |-\vec{b}| = |\vec{b}|)$$

$$|\vec{a} - \vec{b}| \leq |\vec{a}| + |\vec{b}| \text{ (proved)}$$

### Components of vector in 2D

Let  $xoy$  be the co-ordinate plane and  $P(x, y)$  be any point in this plane.

The unit vector along direction of  $x$  axis i.e.  $\vec{ox}$  is denoted by  $\hat{i}$ .

The unit vector along direction of  $y$  axis i.e.  $\vec{oy}$  is denoted by  $\hat{j}$ .


Then from figure - 18 it is clear that  $\vec{om} = x\hat{i}$  and  $\vec{on} = y\hat{j}$

So, the position vector of  $P$  is given

by

$$\vec{op} = \vec{r} = x\hat{i} + y\hat{j}$$

$$\text{And } op = |\vec{op}| = r = \sqrt{x^2 + y^2}$$



### Representation of vector in component form in 2D

If  $\vec{AB}$  is any vector having end points  $A(x_1, y_1)$  and  $B(x_2, y_2)$ , then it can be represented by

$$\vec{AB} = (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j}$$

### Components of Vector in 3D


Let  $P(x, y, z)$  be a point in space and  $\hat{i}, \hat{j}$  and  $\hat{k}$  be the unit vectors along  $x$  axis,  $y$  axis and  $z$  axis respectively. (as shown in fig-19)

Then the position vector of  $P$  is given by

$$\vec{OP} = x\hat{i} + y\hat{j} + z\hat{k}, \text{ the vectors}$$

$x\hat{i}, y\hat{j}, z\hat{k}$  are called the components of  $\vec{OP}$  along  $x$ -axis,  $y$  axis, and  $z$ -axis respectively.

$$\text{And } OP = |\vec{OP}| = \sqrt{x^2 + y^2 + z^2}$$



### Addition and Scalar Multiplication in terms of component form of vectors :-

For any vector  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$

- (i)  $\vec{a} + \vec{b} = (a_1 + b_1)\hat{i} + (a_2 + b_2)\hat{j} + (a_3 + b_3)\hat{k}$
- (ii)  $\vec{a} - \vec{b} = (a_1 - b_1)\hat{i} + (a_2 - b_2)\hat{j} + (a_3 - b_3)\hat{k}$
- (iii)  $k\vec{a} = ka_1\hat{i} + ka_2\hat{j} + ka_3\hat{k}$ , where  $k$  is a scalar.
- (iv)  $\vec{a} = \vec{b} \Leftrightarrow a_1\hat{i} + a_2\hat{j} + a_3\hat{k} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$   
 $\Rightarrow a_1 = b_1, a_2 = b_2, a_3 = b_3$

### Representation of Vector in component form in 3-D and Distance between two points:

If  $\vec{AB}$  is any vector having end points  $A(x_1, y_1, z_1)$  and  $B(x_2, y_2, z_2)$ , then it can be represented by

$$\begin{aligned} \vec{AB} &= \text{Position vector of } B - \text{Position vector of } A \\ &= (x_2\hat{i} + y_2\hat{j} + z_2\hat{k}) - (x_1\hat{i} + y_1\hat{j} + z_1\hat{k}) \\ &= (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j} + (z_2 - z_1)\hat{k} \end{aligned}$$

$$|\vec{AB}| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

### Example 3:-

Show that the points  $A(2, 6, 3)$ ,  $B(1, 2, 7)$  and  $C(3, 10, 1)$  are collinear.

Solution:- From given data position vector of  $A, \vec{OA}$   
 position vector of  $B, \vec{OB} = \hat{i} + 2\hat{j} + 7\hat{k} = 2\hat{i} + 6\hat{j} + 3\hat{k}$   
 position vector of  $C, \vec{OC} = 3\hat{i} + 10\hat{j} - \hat{k}$

$$\text{Now } \vec{AB} = \vec{OB} - \vec{OA} = (1-2)\hat{i} + (2-6)\hat{j} + (7-3)\hat{k} = -\hat{i} - 4\hat{j} + 4\hat{k}$$



Condition of perpendicularity :-

$$\vec{AC} = \vec{OC} - \vec{OA} = (8-2)\hat{i} + (10-6)\hat{j} + (-1-3)\hat{k} = 6\hat{i} + 4\hat{j} - 4\hat{k}$$

$$= -(-6\hat{i} - 4\hat{j} + 4\hat{k}) = -\vec{AB}$$

$\Rightarrow \vec{AB} \parallel \vec{AC}$  or collinear.

$\therefore$  They have same support and common point A.

As 'A' is common to both vector, that proves A, B and C are collinear.

Example-4 - Prove that the points having position vector given by  $2\hat{i} - \hat{j} + \hat{k}$ ,  $\hat{i} - 3\hat{j} - 5\hat{k}$  and  $3\hat{i} - 4\hat{j} - 4\hat{k}$  form a right angled triangle. [2009(W)]

Solution:- Let A, B and C be the vertices of a triangle with position vectors  $2\hat{i} - \hat{j} + \hat{k}$ ,  $\hat{i} - 3\hat{j} - 5\hat{k}$  and  $3\hat{i} - 4\hat{j} - 4\hat{k}$  respectively.

Then,  $\vec{AB} = \text{position vector of B} - \text{position vector of A}$

$$= (1-2)\hat{i} + (-3-(-1))\hat{j} + (-5-1)\hat{k} = -\hat{i} - 2\hat{j} - 6\hat{k}$$

$\vec{BC} = \text{position vector of C} - \text{position vector of B}$

$$= (3-1)\hat{i} + (-4-(-3))\hat{j} + (-4-(-5))\hat{k} = 2\hat{i} - \hat{j} + \hat{k}$$

$\vec{AC} = \text{position vector of C} - \text{position vector of A}$

$$= (3-2)\hat{i} + (-4-(-1))\hat{j} + (-4-1)\hat{k} = \hat{i} - 3\hat{j} - 5\hat{k}$$

Now  $AB = |\vec{AB}| = \sqrt{(-1)^2 + (-2)^2 + (-6)^2} = \sqrt{1+4+36} = \sqrt{41}$

$BC = |\vec{BC}| = \sqrt{2^2 + (-1)^2 + 1^2} = \sqrt{4+1+1} = 6$

$AC = |\vec{AC}| = \sqrt{1^2 + (-3)^2 + (-5)^2} = \sqrt{1+9+25} = \sqrt{35}$

from above  $BC^2 + AC^2 = 6^2 + 35 = 41 = AB^2$

Hence ABC is right angled triangle.

Example-5:- find the unit vector in the direction of the vector  $\vec{a} = 3\hat{i} - 4\hat{j} + \hat{k}$ . (2017-W)

Ans:- The unit vector in the direction of  $\vec{a}$  is given by

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|} = \frac{3\hat{i} - 4\hat{j} + \hat{k}}{\sqrt{3^2 + (-4)^2 + 1^2}} = \frac{3\hat{i} - 4\hat{j} + \hat{k}}{\sqrt{26}} = \frac{3}{\sqrt{26}}\hat{i} - \frac{4}{\sqrt{26}}\hat{j} + \frac{1}{\sqrt{26}}\hat{k}$$

Example-6 find a unit vector in the direction of  $\vec{a} + \vec{b}$  where  $\vec{a} = \hat{i} + \hat{j} - \hat{k}$  and  $\vec{b} = \hat{i} - \hat{j} + 3\hat{k}$ .

Ans:- Let  $\vec{r} = \vec{a} + \vec{b} = (\hat{i} + \hat{j} - \hat{k}) + (\hat{i} - \hat{j} + 3\hat{k}) = 2\hat{i} + 2\hat{k}$

unit vector along direction of  $\vec{a} + \vec{b}$  is given by

$$\hat{r} = \frac{\vec{r}}{|\vec{r}|} = \frac{2\hat{i} + 2\hat{k}}{\sqrt{2^2 + 2^2}} = \frac{2\hat{i} + 2\hat{k}}{\sqrt{8}} = \frac{2}{\sqrt{8}}\hat{i} + \frac{2}{\sqrt{8}}\hat{k}$$

$$= \frac{2}{2\sqrt{2}}\hat{i} + \frac{2}{2\sqrt{2}}\hat{k} = \frac{1}{\sqrt{2}}\hat{i} + \frac{1}{\sqrt{2}}\hat{k}$$

### Angle between the vectors:

As shown in figure-20 angle between two vectors  $\vec{RS}$  and  $\vec{PQ}$  can be determined as follows.

Let  $\vec{OB}$  be a vector parallel to  $\vec{RS}$  and  $\vec{OA}$  is a vector parallel to  $\vec{PQ}$  such that  $\vec{OB}$  and  $\vec{OA}$  intersect each other.

Then  $\theta = \angle AOB =$  angle between  $\vec{RS}$  and  $\vec{PQ}$ .




fig-20

If  $\theta = 0$  then ~~the~~ vectors are said to be parallel.

If  $\theta = \frac{\pi}{2}$  then vectors are said to be orthogonal or perpendicular.

### Dot Product or Scalar Product of Vectors


The scalar product of two vectors  $\vec{a}$  and  $\vec{b}$  whose magnitudes are  $a$  and  $b$  respectively denoted by  $\vec{a} \cdot \vec{b}$  is defined as the scalar  $ab \cos \theta$ , where  $\theta$  is the angle between  $\vec{a}$  and  $\vec{b}$  such that  $0 \leq \theta \leq \pi$

$$\boxed{\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta = ab \cos \theta}$$

### Geometrical meaning of dot product

In figure 21(a),  $\vec{a}$  and  $\vec{b}$  are two vectors having  $\theta$  angle between them. Let M be the foot of the perpendicular drawn from B to OA.

Then OM is the projection of  $\vec{b}$  on  $\vec{a}$  and from figure-21(a) it is clear



that,

$$|OM| = |OB| \cos \theta = |\vec{b}| \cos \theta.$$

Now  $\vec{a} \cdot \vec{b} = |\vec{a}| (|\vec{b}| \cos \theta) = |\vec{a}| \times \text{projection of } \vec{b} \text{ on } \vec{a}$   
which gives projection of  $\vec{b}$  on  $\vec{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|}$

Similarly we can write  $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$

$$= |\vec{b}| (|\vec{a}| \cos \theta) = |\vec{b}| \times \text{projection of } \vec{a} \text{ on } \vec{b}$$



Similarly, let us draw a perpendicular from A on OB and let N be the foot of the perpendicular in Fig-21(b).

Then ON = projection of  $\vec{a}$  on  $\vec{b}$   
and  $ON = OA \cos \theta = |\vec{a}| \cos \theta$




Fig-21(b)

### Properties of Dot Product

(i)  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$  (commutative)

(ii)  $\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$  (distributive)

(iii) If  $\vec{a} \parallel \vec{b}$ , then  $\vec{a} \cdot \vec{b} = ab$  { as  $\theta = 0$  in this case  $\cos 0 = 1$  }  
in particular  $(\vec{a})^2 = \vec{a} \cdot \vec{a} = |\vec{a}|^2$   
 $\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1$

(iv) If  $\vec{a} \perp \vec{b}$  then  $\vec{a} \cdot \vec{b} = 0$ . { as  $\theta = 90^\circ$  in this case  $\cos 90^\circ = 0$  }

in particular  $\hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0 = \hat{j} \cdot \hat{i} = \hat{k} \cdot \hat{j} = \hat{i} \cdot \hat{k}$

(v)  $\vec{a} \cdot \vec{0} = \vec{0} \cdot \vec{a} = 0$

(vi)  $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = |\vec{a}|^2 - |\vec{b}|^2 = a^2 - b^2$  { where  $|\vec{a}| = a$  and  $|\vec{b}| = b$  }

(vii) work done by force: - The work done by a force  $\vec{F}$  acting on a body causing displacement  $\vec{d}$  is given by  $w = \vec{F} \cdot \vec{d}$ .

Dot product in terms of rectangular components

For any vectors  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$

we have,

$$\vec{a} \cdot \vec{b} = a_1b_1 + a_2b_2 + a_3b_3 \quad \text{(by applying distributive (i), (ii) and (iv) successively)}$$

Angle between two non zero vectors

For any two non zero vectors  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and

$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ , having  $\theta$  is the angle between

them we have

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{ab} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{a_1b_1 + a_2b_2 + a_3b_3}{\sqrt{a_1^2 + a_2^2 + a_3^2} \sqrt{b_1^2 + b_2^2 + b_3^2}}$$

(in terms of

$$\theta = \cos^{-1} \left( \frac{a_1b_1 + a_2b_2 + a_3b_3}{\sqrt{a_1^2 + a_2^2 + a_3^2} \sqrt{b_1^2 + b_2^2 + b_3^2}} \right) \quad \text{components}$$

Condition of perpendicularity:

Two vectors  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$  are perpendicular to each other  $\Rightarrow a_1b_1 + a_2b_2 + a_3b_3 = 0$

Condition of parallelism:

Two vectors  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$  are parallel to each other  $\Rightarrow \frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$

Section 2 Vector projections of two vectors (Important formulae)

Scalar projection of  $\vec{b}$  on  $\vec{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|}$

Vector projection of  $\vec{b}$  on  $\vec{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|^2} \vec{a} = \left[ \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|^2} \right] \vec{a}$

Scalar projection of  $\vec{a}$  on  $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$

Vector projection of  $\vec{a}$  on  $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \vec{b}$

Examples:—

Q.7 Find the value of  $p$  for which the vectors

$3\hat{i} + 2\hat{j} + 9\hat{k}$ ,  $\hat{i} + p\hat{j} + 3\hat{k}$  are perpendicular to each other.

Solution :- Let  $\vec{a} = 3\hat{i} + 2\hat{j} + 9\hat{k}$  and  $\vec{b} = \hat{i} + p\hat{j} + 3\hat{k}$ .

Here  $a_1 = 3, a_2 = 2, a_3 = 9$

$b_1 = 1, b_2 = p$  &  $b_3 = 3$

Given  $\vec{a} \perp \vec{b} \Rightarrow a_1b_1 + a_2b_2 + a_3b_3 = 0$

$$\Rightarrow 3 \cdot 1 + 2 \cdot p + 9 \cdot 3 = 0$$



$$\Rightarrow 3 + 2p + 27 = 0$$

$$\Rightarrow 3 + 2p + 27 = 0$$

$$\Rightarrow 2p = -30$$

$$\Rightarrow p = -15 \text{ (Ans)}$$

Q.8 Find the value of  $p$  for which the vectors  $\vec{a} = 3\hat{i} + 2\hat{j} + 9\hat{k}$ ,  $\vec{b} = \hat{i} + p\hat{j} + 3\hat{k}$  are parallel to each other. (2014-W)

Solution :- Given  $\vec{a} \parallel \vec{b} \Leftrightarrow \frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3} \Leftrightarrow \frac{3}{1} = \frac{2}{p} = \frac{9}{3}$

$\Leftrightarrow 3 = \frac{2}{p} \Leftrightarrow p = \frac{2}{3}$  (Ans) {Taking 1<sup>st</sup> two terms}

{Note - any two expression may be taken for finding  $p$ .}

Q.9 Find the scalar product of  $3\hat{i} - 4\hat{j}$  and  $-2\hat{i} + \hat{j}$ . (2015-5)

Solution :-  $(3\hat{i} - 4\hat{j}) \cdot (-2\hat{i} + \hat{j}) = (3 \times (-2)) + ((-4) \times 1) = (-6) + (-4) = -10$

Q.10 Find the angle between the vectors  $5\hat{i} + 3\hat{j} + 4\hat{k}$  and  $6\hat{i} - 8\hat{j} - \hat{k}$ . (2015-W)

Solution :- Let  $\vec{a} = 5\hat{i} + 3\hat{j} + 4\hat{k}$  and  $\vec{b} = 6\hat{i} - 8\hat{j} - \hat{k}$   
Let  $\theta$  be the angle between  $\vec{a}$  and  $\vec{b}$ .

$$\text{Then } \theta = \cos^{-1} \left( \frac{a_1 b_1 + a_2 b_2 + a_3 b_3}{\sqrt{a_1^2 + a_2^2 + a_3^2} \sqrt{b_1^2 + b_2^2 + b_3^2}} \right)$$

$$= \cos^{-1} \left( \frac{5 \cdot 6 + 3 \cdot (-8) + 4 \cdot (-1)}{\sqrt{5^2 + 3^2 + 4^2} \sqrt{6^2 + (-8)^2 + (-1)^2}} \right) = \cos^{-1} \left( \frac{30 - 24 - 4}{\sqrt{50} \sqrt{101}} \right)$$

Q11. Find the Scalar and vector projection of  $\vec{a}$  on  $\vec{b}$   
where  $\vec{a} = \hat{i} - \hat{j} - \hat{k}$  and  $\vec{b} = 3\hat{i} + \hat{j} + 3\hat{k}$  (2013-w, 2017-w)

Solution: Scalar Projection of  $\vec{a}$  on  $\vec{b} =$

$$\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{1 \cdot 3 + (-1) \cdot 1 + (-1) \cdot 3}{\sqrt{3^2 + 1^2 + 3^2}} = \frac{3-1-3}{\sqrt{19}} = \frac{-1}{\sqrt{19}}$$

Vector projection of  $\vec{a}$  on  $\vec{b}$ .

$$\begin{aligned} &= \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \vec{b} = \frac{1 \cdot 3 + (-1) \cdot 1 + (-1) \cdot 3}{(\sqrt{3^2 + 1^2 + 3^2})^2} (3\hat{i} + \hat{j} + 3\hat{k}) \\ &= \frac{3-1-3}{19} (3\hat{i} + \hat{j} + 3\hat{k}) = \frac{-1}{19} (3\hat{i} + \hat{j} + 3\hat{k}) \end{aligned}$$

Q12. Find the Scalar and Vector projection of  $\vec{b}$  on  $\vec{a}$   
where  $\vec{a} = 3\hat{i} + \hat{j} - 2\hat{k}$  and  $\vec{b} = 2\hat{i} + 3\hat{j} - 4\hat{k}$  {2015-S}

Solution: Scalar Projection of  $\vec{b}$  on  $\vec{a}$

$$= \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|} = \frac{3 \cdot 2 + 1 \cdot 3 + (-2) \cdot (-4)}{(\sqrt{3^2 + 1^2 + (-2)^2})^2} (3\hat{i} + \hat{j} - 2\hat{k})$$

$$= \frac{17}{14} (3\hat{i} + \hat{j} - 2\hat{k})$$



Q-13 If  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$ , then prove that  $\vec{a} = \vec{0}$  or  $\vec{b} = \vec{c}$  or  $\vec{a} \perp (\vec{b} - \vec{c})$  Proof :- Given  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$

$$\Rightarrow (\vec{a} \cdot \vec{b}) - (\vec{a} \cdot \vec{c}) = 0$$

$$\Rightarrow \vec{a} \cdot (\vec{b} - \vec{c}) = 0 \quad \left( \begin{array}{l} \text{applying} \\ \text{distributive property} \end{array} \right)$$

Dot product of above two vectors is zero indicates the following conditions

$$\vec{a} = \vec{0} \text{ or } \vec{b} - \vec{c} = \vec{0} \text{ or } \vec{a} \perp (\vec{b} - \vec{c})$$

$$\Rightarrow \vec{a} = \vec{0} \text{ or } \vec{b} = \vec{c} \text{ or } \vec{a} \perp (\vec{b} - \vec{c}) \text{ (proved)}$$

Example -14 Find the work done by force  $\vec{F} = \hat{i} + \hat{j} - \hat{k}$  acting on a particle if the particle is displaced from A(3,3,3) to B(4,4,4)

Ans :- Let O be the origin, then

$$\text{Position vector of A } \vec{OA} = 3\hat{i} + 3\hat{j} + 3\hat{k}$$

$$\text{Position vector of B } \vec{OB} = 4\hat{i} + 4\hat{j} + 4\hat{k}$$

## Limit and continuity

In Mathematics, Differential calculus is a subfield of calculus that studies the ratio at which quantities change.

### Quantity

A quantity is an amount, number or measurement, in other words an expression having value considered as a whole. These numbers can be expressed as a whole number, fraction, Decimals, Percentages.

~~and~~

### Type of Quantities

Quantities are broadly divided

in to two categories.

- (1) Constant
- (2) variable



## Constant

Constant is a symbol which remains the same value through out a set of mathematical operations. There are mainly two types of Constant.

- (i) absolute constant
- (ii), arbitrary constant.

### (i) Absolute Constant :-

Operation we may perform on absolute constants do not change whatever known.

### (ii) Arbitrary Constant :-

In the equation  $y = ax + b$  of a straight line are arbitrary

constants as they remain fixed for a particular straight line but vary from straight line where  $a$  &  $b$  are arbitrary constants.

## (2) variable

Variable is a symbol which can take various numerical values. Variables are of two types.

- (i). Dependent variable
- (ii). Independent variable

Ex:  $y = 2x + 4$

when  $x = 1$  then  $y = 6$ , The value of  $y$  keeps on changing dependently upon the value of  $x$ . So  $x$  is the independent variable and  $y$  is the dependent variable.



## Set

A set is the collection of any well defined objects known as elements or numbers of the set.

$$\text{Ex: } A = \{1, 2, 3, 4, 5\}$$

## Relation

A relation between two sets is a collection of ordered pairs containing one object from each set. If the object  $x$  is from 1st set and the object  $y$  is from 2nd set then the objects are said to be related if the ordered pair  $(x, y)$  is in the relation.

Ex:

$$A = \{1, 2\}$$

$$B = \{1, 2, 3\}$$

$$A \times B = \{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3)\}$$

↳ Cartesian Product.

(i). find a relation where  
 $x = y \rightarrow R_1 = \{(1,1), (2,2)\}$

(ii) find a Relation where  $x = y^2$   
 $\rightarrow R_2 = \{(1,1)\}$

(iii) find a relation where  $x^2 = y \rightarrow R_3 = \{(1,1)\}$


## function

A function is a special case of relation. Let  $(x,y)$  be two nonempty sets and  $R$  be a relation from  $x$  to  $y$ . then  $R$  may not relate an element of  $x$  to an element of  $y$  or it may relate an element of  $x$  to more than one element of  $y$ .

But a function relates each element of  $x$  to a unique element of  $y$   
 $f: x \rightarrow y$  (reads as " $f$  maps  $x$  to  $y$ ")



Pictorially a function can be represented as



### Main features of function:—

- (i). To each element  $x \in X$ , there exists a unique element  $y \in Y$  such that  $y = f(x)$
- (ii). Distinct elements of  $X$  may be associated with the same elements of  $Y$ .
- (iii). These may be elements of  $Y$  which are not associated with any element of  $X$ .

## Domain

The set  $x$  is called the Domain of the function  $f$ .


## Range

The set of all images of the elements of  $x$  under the mapping  $f$  is called the range of  $f$  is denoted by  $f(x)$ .  
in general  $f(x) \subseteq y$ .


## Intervals

Let 'a' & 'b' two distinct real numbers  
and set  $a < b$ .  
Then (i).  $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$  is called  
Closed interval from a to b including

a & b.




(ii)  $(a, b) = \{x \in \mathbb{R} : a < x < b\}$  is called  
Open interval from a to b excluding a & b.  
(not included) (not included)






(iii).  $[a, b) = \{x \in \mathbb{R} : a \leq x < b\}$  is called  
Semi closed and semi open interval.



(iv)  $(a, b] = \{x \in \mathbb{R} : a < x \leq b\}$  is called  
Semi open and semi closed interval.



Notes

$\infty$  will always comes with  
open Brackett, because we can't show  
 $\infty$  on a number line. (as it is undefined)

(v).  $[a, \infty) \rightarrow \{x \in \mathbb{R} : x \geq a\}$

(vi).  $(a, \infty) \rightarrow \{x \in \mathbb{R} : x > a\}$

(vii).  $(-\infty, a] \rightarrow \{x \in \mathbb{R} : x \leq a\}$


(viii).  $(-\infty, a) \rightarrow \{x \in \mathbb{R} : x < a\}$

(ix).  $(-\infty, \infty) \rightarrow \{x \in \mathbb{R}\}$

## Classification of function

A function  $f$  defined from the set  $X$  to the set  $Y$  is said to be an onto function if range of  $f$  is  $Y$ , i.e. a Proper Subset of  $f$  in  $Y$ .

on  
The function  $f: X \rightarrow Y$  is called an onto function if there exist at least one element of  $Y$  which does not correspond to any element of  $X$ .



## onto function (surjective mapping):

A function  $f: X \rightarrow Y$  is said to be an onto function if every element  $y$  in  $Y$  is the image of some element in  $X$ .




$$\text{If } X = \{1, 2, 3, 4\}$$

$$\text{or } Y = \{a, b, c\}$$


$$F = \{(1, a), (2, b), (3, c), (4, c)\}$$

(Y set is completely used)



One one mapping


If distinct element of X have distinct image in Y then the function called one one function.



One one and onto function (Bijection)


A function which is such that it is (i) onto (ii) one one is called Bijection.

$$X = \{1, 2, 3\}; Y = \{a, b, c\}$$



## Many-one function :-

A function  $f$  from the set  $X$  to the set  $Y$  is said to be ~~one~~ many one if there exists at least one element in  $Y$  which has more than one preimage in  $X$ .





## ⇒ Types of Function :-


### ① Identity Function :-

Def<sup>n</sup> :- The Function  $f$  defined by  $f(x) = x$  for  $\forall x \in \mathbb{R}$  is called the identity Function.

$$\text{Domain} = \mathbb{R}$$

$$\text{Range} = \mathbb{R}$$

Graph :-




### ② Constant Function :-

Def<sup>n</sup> :- The Function  $f$  defined by  $f(x) = \frac{1}{x}$  is called Reciprocal Function.

$$\text{Domain} = \mathbb{R}$$

$$\text{Range} = \mathbb{C}$$

Graph :-




### ③ Reciprocal Function :-

Def<sup>n</sup> :- The Function  $f$  defined by  $f(x) = \frac{1}{x}$  is called Reciprocal Function.

$$\text{Domain} = \mathbb{R} - \{0\}$$

$$\text{Range} = \mathbb{R} - \{0\}$$

Graph :-



### ④ Modulus Function :-


Def<sup>n</sup> :- The Function  $f$  defined by  $f(x)$ .

$$f(x) = |x| = \begin{cases} x & \text{when } x \geq 0 \\ -x & \text{when } x < 0 \end{cases}$$

is called the modulus or absolute value Function.

$$\text{Domain} = \mathbb{R}, \text{Range} = [0, \infty)$$

Graph :-



### ⑤ Greatest integer Function :-

Def<sup>n</sup> :- The Function  $f$  defined by  $f(x) = [x]$  For all  $x \in \mathbb{R}$  is called the greatest integer Function.

where  $[x] = k$  iff  $k \leq x < k+1$

Domain =  $\mathbb{R}$


Range =  $\mathbb{Z}$

$$[n+] = n$$

$$[n-] = n-1$$

$$[n] = n$$

Graph :-



### ⑥ Signum Function :-

Def<sup>n</sup> :- The Function  $f$  defined by  $f(x) = \begin{cases} \frac{|x|}{x} & , x \neq 0 \\ 0 & , \text{when } x = 0 \end{cases}$


in other words

$$f(x) = \begin{cases} 1 & x > 0 \\ 0 & x = 0 \\ -1 & x < 0 \end{cases}$$

Domain =  $\mathbb{R}$

Range =  $\{1, 0, -1\}$

Graph :-





### ⑦ Logarithmic Function :-

Def<sup>n</sup> :- The Function  $f$  defined by  $f(x) = \log_a x$  where  $x > 0$ ,  $a > 0$  &  $a \neq 1$  is called the Logarithmic Function.

Domain =  $(0, \infty)$  , Range =  $\mathbb{R}$

Graph :-





## properties of Logarithmic Function :-

$$(i) \log mn = \log m + \log n$$


$$(ii) \log m^n = n \log m$$

$$(iii) \log \frac{m}{n} = \log m - \log n$$

## Exponential Function :-

The function  $f$  defined by  $f(x) = a^x$ , where  $a > 0$  and  $x \in \mathbb{R}$  is called exponential function.

$$D = \mathbb{R}, \quad R = (0, \infty)$$



## Rational Function :-

The function  $f$  defined by  $f(x) = \frac{g(x)}{h(x)}$ , where  $g(x)$  and  $h(x)$  are polynomial functions and  $h(x) \neq 0$  is called a Rational Function.

Domain :- Real no set except those values of  $x$  for which  $h(x) = 0$


## Trigonometric Function :- (Circular Function)

$\sin x$ ,  $\cos x$ ,  $\tan x$ ,  $\cot x$ ,  $\sec x$ ,  $\csc x$  are trigonometric functions.

(i) Sine Function :-  $f(x) = \sin x$

Domain =  $\mathbb{R}$

Range =  $[-1, 1]$




## Square root Function :-

The function  $f$  defined by  $f(x) = \sqrt{x}$  is called the square root function.

Domain =  $[0, \infty)$

Range =  $[0, \infty)$

Graph :-



## Polynomial Function :-

The function  $f$  defined by  $f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$   
 $a_n \neq 0$ , where  $a_0, a_1, a_2, \dots, a_n$  are real nos and  $n \in \mathbb{N}$  is  
called a polynomial function of degree  $n$ .

Domain =  $\mathbb{R}$ , Range =  $\mathbb{R}$

## $\Rightarrow$ Inverse Trigonometry :-

| Function Name                | Domain                           | Range                     | Graph |
|------------------------------|----------------------------------|---------------------------|-------|
| $\sin^{-1}x$                 | $[-1, 1]$                        | $[-\pi/2, \pi/2]$         |       |
| $\cos^{-1}x$                 | $[-1, 1]$                        | $[0, \pi]$                |       |
| $\tan^{-1}x$                 | $(-\infty, \infty)$              | $(-\pi/2, \pi/2)$         |       |
| $\cot^{-1}x$                 | $(-\infty, \infty)$              | $(0, \pi)$                |       |
| $\sec^{-1}x$                 | $(-\infty, -1] \cup [1, \infty)$ | $[0, \pi] - \{\pi/2\}$    |       |
| $\operatorname{cosec}^{-1}x$ | $(-\infty, -1] \cup [1, \infty)$ | $[-\pi/2, \pi/2] - \{0\}$ |       |




Even Function :- A Function  $f(x)$  is said to be an even Function if  $f(-x) = f(x)$

odd Function :- A Function  $f(x)$  is said to be an odd Function if  $f(-x) = -f(x)$

cosine Function :-  $f(x) = \cos x$

$$\text{Domain} = \mathbb{R}$$


$$\text{Range} = [-1, 1]$$



Tangent Function :-  $f(x) = \tan x$

$$\text{Domain} = \mathbb{R} - \{(2k+1)\pi/2 : k \in \mathbb{Z}\}$$


$$\text{Range} = \mathbb{R}$$



secant Function :-  $f(x) = \sec x$

$$\text{Domain} = \mathbb{R} - \{(2n+1)\pi/2 : n \in \mathbb{Z}\}$$


$$\text{Range} = \mathbb{R} - (-1, 1)$$



cotangent Function :-  $f(x) = \cot x$

$$\text{Domain} = \mathbb{R} - n\pi \quad n \in \mathbb{Z}$$


$$\text{Range} = \mathbb{R}$$



cosecant Function :-  $f(x) = \csc x$

$$\text{Domain} = \mathbb{R} - n\pi$$

$$\text{Range} = \mathbb{R} - (-1, 1)$$



Explicitly Function :-

A Function which is expressed directly in terms of independence variable is called an explicit Function.  $y = x^2 - 2x + 5$

Implicit Function :-

If a function is not expressed directly in terms of independent variable is called an implicit Function.

Single valued Function :-

A Function  $y = f(x)$  is said to be a single valued Function if there is one and only value of  $y$  corresponding to each value of  $x$ .



## Periodic Function :-

A function  $f(x)$  is said to be a periodic function if there exists a positive real constant  $T$  such that  $f(x+T) = f(x)$ , for all  $x \in \mathbb{R}$ .

## Introduction to Limits :-

The notion of closeness and nearness is basic in several branches of Mathematics. The concept of limit of a function, which is fundamental in calculus, is based on this notion.

consider the function,

$$f: \mathbb{R} \rightarrow \mathbb{R} \text{ defined by}$$

$f(x) = 2x+1$ . Let the variable  $x$  takes values closer and closer to 2.

Table-1

|        |     |      |       |        |
|--------|-----|------|-------|--------|
| $x$    | 2.1 | 2.01 | 2.001 | 2.0001 |
| $f(x)$ | 5.2 | 5.02 | 5.002 | 5.0002 |

Table-2

|        |     |      |       |        |
|--------|-----|------|-------|--------|
| $x$    | 1.9 | 1.99 | 1.999 | 1.9999 |
| $f(x)$ | 4.8 | 4.98 | 4.998 | 4.9998 |

Symbolically,

$$\lim_{x \rightarrow 2} (2x+1) = 5$$

$$x \rightarrow 2$$

From table-2, it is observed that the difference between  $x$  and 2 is decreasing, the difference between  $f(x)$  and 5 is also decreasing correspondingly.

## Limit of a function :-


Def<sup>n</sup>:- Let  $f(x)$  be a function defined in some neighbourhood of  $a$ , except possibly at  $a$  and  $L$  be a number, we say that limit of  $f(x)$  as  $x$  approaches  $a$  is ' $L$ ' written  $\lim_{x \rightarrow a} f(x) = L$ . If for any  $\epsilon > 0$  however small there exists  $\delta$ ,  $\delta > 0$  such that

$$|f(x) - L| < \epsilon, \text{ whenever } 0 < |x - a| < \delta$$

\* To each  $\epsilon > 0$ , there exists a positive no.  $\delta$  such that,

$$\text{when } 0 < |x - a| < \delta \Rightarrow |f(x) - L| < \epsilon$$





(i)  $(\epsilon - \delta)$  method :-

Ex-1 Show that  $\lim_{x \rightarrow 1} (3x+2) = 5$

Solution -  $\lim_{x \rightarrow a} f(x) = L$

$f(x) = 3x+2$  ,  $L = 5$  ,  $a = 1$

$$|f(x) - L| < \epsilon$$

$$|(3x+2) - 5| < \epsilon$$

$$|3x - 3| < \epsilon$$

$$3|x-1| < \epsilon$$

$$|x-1| < \epsilon/3$$

$$\boxed{|x-1| < \delta}$$

$(\because \text{choose } \frac{\epsilon}{3} = \delta)$

so,  $|(3x+2) - 5| < \epsilon \Rightarrow |x-1| < \delta$

Hence  $\lim_{x \rightarrow 1} (3x+2) = 5$

Evaluation of Left hand and Right hand Limit :-

Left hand Limit :- To evaluate L.H.L of  $f(x)$  at  $x=a$

$$\lim_{x \rightarrow a^-} f(x)$$

Step-1 :- write  $\lim_{x \rightarrow a^-} f(x)$

Step-2 :- put  $x = a-h$  and replace  $x \rightarrow a-h$  by  $h \rightarrow 0$  to obtain

$$\lim_{h \rightarrow 0} f(a-h)$$

Step-3 :- simplify  $\lim_{h \rightarrow 0} f(a-h)$  by using the

$\epsilon$ - $\delta$  method :-

$$|f(x) - L| < \epsilon \text{ whenever } |x - a| < \delta$$

→ prove that  $\lim_{x \rightarrow 2} (2x+1) = 5$  using the  $\epsilon$ - $\delta$  method / definition method?

Ans.  $f(x) = 2x+1$ ,  $L = 5$ ,  $a = 2$

using the definition of limit  $|f(x) - L| < \epsilon$

$$|(2x+1) - 5| < \epsilon$$

$$\Rightarrow |2x - 4| < \epsilon$$

$$\Rightarrow |2(x-2)| < \epsilon$$

$$\Rightarrow |x-2| < \epsilon/2$$

$$\Rightarrow |x-2| < \delta \quad (\text{let, } \epsilon/2 = \delta)$$

$$\text{So, } |(2x+1) - 5| < \epsilon = |x-2| < \delta \quad (\text{proved})$$

→  $\lim_{x \rightarrow 3} (3x+4) = 13$

Ans.  $f(x) = 3x+4$ ,  $L = 13$ ,  $a = 3$

using the definition of limit  $|f(x) - L| < \epsilon$

$$|(3x+4) - 13| < \epsilon$$

$$\Rightarrow |3x - 9| < \epsilon$$

$$\Rightarrow |3(x-3)| < \epsilon$$

$$\Rightarrow |x-3| < \epsilon/3$$

$$\Rightarrow |x-3| < \delta \quad (\text{let, } \epsilon/3 = \delta)$$

$$\text{So, } |(3x+4) - 13| < \epsilon = |x-3| < \delta \quad (\text{proved})$$

→  $\lim_{x \rightarrow -1} (4x-5) = -9$

Ans.  $f(x) = 4x-5$ ,  $L = -9$ ,  $a = -1$

$$|(4x-5) - (-9)| < \epsilon$$

$$\Rightarrow |4x - 5 + 9| < \epsilon$$



$$\Rightarrow |4x+4| < \epsilon$$

$$\Rightarrow |4(x+1)| < \epsilon$$

$$\Rightarrow |x+1| < \epsilon/4$$

$$\Rightarrow |x+1| < \delta \quad (\text{Let, } \epsilon/4 = \delta)$$

$$\text{so, } |(4x-5) - (-9)| < \epsilon = |x - (-1)| < \delta$$

$$\Rightarrow |(4x-5) + 9| < \epsilon = |x+1| < \delta \quad (\text{proved})$$

$\Rightarrow$  L.H.L and R.H.L

L.H.L

$$\lim_{x \rightarrow c^-} f(x)$$

R.H.L

$$\lim_{x \rightarrow c^+} f(x)$$

Ex-1

check the existence of the function  $f(x) = 3x+4$  at  $x=2$ .

Ans.

| L.H.L                                                                                                            | R.H.L                                                                                                            |
|------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|
| $\lim_{x \rightarrow c^-} f(x)$ $= \lim_{x \rightarrow 2^-} (3x+4)$ $= \lim_{h \rightarrow 0} (3(2-h)+4)$ $= 10$ | $\lim_{x \rightarrow c^+} f(x)$ $= \lim_{x \rightarrow 2^+} (3x+4)$ $= \lim_{h \rightarrow 0} (3(2+h)+4)$ $= 10$ |

$\therefore$  L.H.L = R.H.L the limiting value exists.

Ex-2

$f(x) = [x]$  at  $x=3$

Ans.

| L.H.L                                                                                                   | R.H.L                                                                                                   |
|---------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|
| $\lim_{x \rightarrow c^-} f(x)$ $= \lim_{x \rightarrow 3^-} [x]$ $= \lim_{h \rightarrow 0} [3-h]$ $= 2$ | $\lim_{x \rightarrow c^+} f(x)$ $= \lim_{x \rightarrow 3^+} [x]$ $= \lim_{h \rightarrow 0} [3+h]$ $= 3$ |

## → Evaluation of Limit :-

- ① Algebraic problem
- ② Trigonometric problem
- ③ Exponential or logarithmic

→ Algebraic method

- Direct substitution method
- Factorisation method
- Rationalisation method
- some standard formula
- when  $x \rightarrow \infty$  (infinity)

### Direct substitution method :-

- ①  $\lim_{x \rightarrow 1} (2x + 3) = 2 + 3 = 5$
- ②  $\lim_{x \rightarrow 2} \left( \frac{4x+2}{3x-1} \right) = \frac{10}{5} = 2$
- ③  $\lim_{x \rightarrow 0} (4x^5 - 3x^3 + 2x^2 + 1) = 1$
- ④  $\lim_{x \rightarrow -1} (3x-2)(4x-3) = -5 \times -7 = 35$

### Factorisation method :-

- ①  $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} \left( \frac{0}{0} \text{ form} \right)$   
 $= \lim_{x \rightarrow 2} \frac{(x-2)(x+2)}{(x-2)}$   
 $= \lim_{x \rightarrow 2} x + 2 = 2 + 2 = 4$
- ②  $\lim_{x \rightarrow 1} \frac{x^2 - 2x + 1}{x - 1} \left( \frac{0}{0} \text{ form} \right)$   
 $= \lim_{x \rightarrow 1} \frac{(x-1)^2}{(x-1)}$   
 $= \lim_{x \rightarrow 1} x - 1 = 1 - 1 = 0$
- ③  $\lim_{x \rightarrow 0} \frac{x^2 - x}{x} \left( \frac{0}{0} \text{ form} \right)$   
 $= \lim_{x \rightarrow 0} \frac{x(x-1)}{x} = 0 - 1 = -1$



### Rationalisation method :-

$$\textcircled{1} \lim_{x \rightarrow 0} \frac{x}{\sqrt{x+1} - 1} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1} + 1)}{(\sqrt{x+1} - 1)(\sqrt{x+1} + 1)}$$

$$= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1} + 1)}{(\sqrt{x+1})^2 - (1)^2}$$

$$= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1} + 1)}{x + 1 - 1}$$

$$= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1} + 1)}{x} = \sqrt{1} + 1 = 1 + 1 = 2$$

$$\textcircled{2} \lim_{x \rightarrow 0} \frac{\sqrt{x^2+4} - 2}{x^2} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 0} \frac{(\sqrt{x^2+4} - 2)(\sqrt{x^2+4} + 2)}{x^2(\sqrt{x^2+4} + 2)}$$

$$= \lim_{x \rightarrow 0} \frac{(\sqrt{x^2+4})^2 - (2)^2}{x^2(\sqrt{x^2+4} + 2)}$$

$$= \lim_{x \rightarrow 0} \frac{x^2 + 4 - 4}{x^2(\sqrt{x^2+4} + 2)}$$

$$= \lim_{x \rightarrow 0} \frac{x^2}{x^2(\sqrt{x^2+4} + 2)} = \frac{1}{\sqrt{4} + 2} = \frac{1}{2+2} = \frac{1}{4}$$

### using standard formula :-

using standard formula =

$$\boxed{\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1}}$$

$$\textcircled{1} \lim_{x \rightarrow 4} \frac{x^2 - 16}{x - 4} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 4} \frac{(x)^2 - (4)^2}{x - 4} = 2 \times 4^{2-1} = 2 \times 4 = 8$$

$$\textcircled{2} \lim_{x \rightarrow 2} \frac{x^3 - 8}{x - 2} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 2} \frac{(x)^3 - (2)^3}{x - 2} = 3 \times 2^{3-1} = 3 \times 2^2 = 12$$

### Rationalisation method :-

$$\textcircled{1} \lim_{x \rightarrow 0} \frac{x}{\sqrt{x+1} - 1} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1} + 1)}{(\sqrt{x+1} - 1)(\sqrt{x+1} + 1)}$$

$$= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1} + 1)}{(\sqrt{x+1})^2 - (1)^2}$$

$$= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1} + 1)}{x + 1 - 1}$$

$$= \lim_{x \rightarrow 0} \frac{x(\sqrt{x+1} + 1)}{x} = \sqrt{1} + 1 = 1 + 1 = 2$$

$$\textcircled{2} \lim_{x \rightarrow 0} \frac{\sqrt{x^2+4} - 2}{x^2} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 0} \frac{(\sqrt{x^2+4} - 2)(\sqrt{x^2+4} + 2)}{x^2(\sqrt{x^2+4} + 2)}$$

$$= \lim_{x \rightarrow 0} \frac{(\sqrt{x^2+4})^2 - (2)^2}{x^2(\sqrt{x^2+4} + 2)}$$

$$= \lim_{x \rightarrow 0} \frac{x^2 + 4 - 4}{x^2(\sqrt{x^2+4} + 2)}$$

$$= \lim_{x \rightarrow 0} \frac{x^2}{x^2(\sqrt{x^2+4} + 2)} = \frac{1}{\sqrt{4} + 2} = \frac{1}{2+2} = \frac{1}{4}$$

using standard formula:-

using standard formula =

$$\boxed{\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1}}$$

$$\textcircled{1} \lim_{x \rightarrow 4} \frac{x^2 - 16}{x - 4} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 4} \frac{(x)^2 - (4)^2}{x - 4} = 2 \times 4^{2-1} = 2 \times 4 = 8$$

$$\textcircled{2} \lim_{x \rightarrow 2} \frac{x^3 - 8}{x - 2} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 2} \frac{(x)^3 - (2)^3}{x - 2} = 3 \times 2^{3-1} = 3 \times 2^2 = 12$$



$$\textcircled{3} \quad \lim_{x \rightarrow 5} \frac{\sqrt{x} - \sqrt{5}}{x - 5} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 5} \frac{(x)^{1/2} - (5)^{1/2}}{x - 5} = \frac{1}{2} \times 5^{\frac{1}{2}-1} = \frac{1}{2} \times 5^{-1/2} = \frac{1}{2} \times \frac{1}{\sqrt{5}} = \frac{1}{2\sqrt{5}}$$

$$\textcircled{4} \quad \lim_{x \rightarrow 1} \frac{x^5 - 1}{x - 1} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 1} \frac{(x)^5 - (1)^5}{x - 1} = 5 \times 1^{5-1} = 5 \times 1 = 5$$

$$\textcircled{5} \quad \lim_{x \rightarrow -1} \frac{x^3 + 1}{x + 1} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow -1} \frac{x^3 - (-1)}{x - (-1)}$$

$$= \lim_{x \rightarrow -1} \frac{(x)^3 - (-1)^3}{x - (-1)} = 3 \times (-1)^{3-1} = 3 \times (-1)^2 = 3 \times 1 = 3$$

$$\textcircled{6} \quad \lim_{x \rightarrow 2} \frac{x^5 - 32}{x^3 - 8} \quad \left(\frac{0}{0} \text{ form}\right)$$

$$= \lim_{x \rightarrow 2} \frac{(x)^5 - (2)^5}{(x)^3 - (2)^3}$$

Divide the numerator and denominator by  $(x-2)$ .

$$= \lim_{x \rightarrow 2} \frac{\frac{(x)^5 - (2)^5}{x - 2}}{\frac{(x)^3 - (2)^3}{x - 2}}$$

$$= \lim_{x \rightarrow 2} \frac{(x)^5 - (2)^5}{x - 2} \div \lim_{x \rightarrow 2} \frac{(x)^3 - (2)^3}{x - 2} = \frac{5 \times (2)^{5-1}}{3 \times (2)^{3-1}} = \frac{5 \times 2^4}{3 \times 2^2} = \frac{80}{12}$$

$$\textcircled{7} \quad \lim_{x \rightarrow 0} \frac{(3+2x)^3 - 27}{x} \quad \left(\frac{0}{0} \text{ form}\right)$$

Ans  $\lim_{x \rightarrow 0} \frac{(3+2x)^3 - (3)^3}{x}$

$$= \lim_{x \rightarrow 0} \frac{(3+2x)^3 - (3)^3}{3+2x - 3}$$

$$= \lim_{3+2x \rightarrow 3} \frac{(3+2x)^3 - (3)^3}{(3+2x) - 3} = 2 \times 3 \times (3)^{3-1} = 2 \times 3 \times 3^2 = 54$$

$$\textcircled{8} \quad \lim_{x \rightarrow 0} \frac{(4-5x)^3 - 64}{x}$$

$$\text{Ans.} \quad \lim_{x \rightarrow 0} \frac{(4-5x)^3 - (4)^3}{x}$$

$$-5 \quad \lim_{x \rightarrow 0} \frac{(4-5x)^3 - (4)^3}{4-5x-4}$$

$$-5 \quad \lim_{4-5x \rightarrow 4} \frac{(4-5x)^3 - (4)^3}{(4-5x) - 4} = -5 \times 3 \times (4)^{3-1} = -5 \times 3 \times 4^2 = -240$$

$$\textcircled{9} \quad \lim_{x \rightarrow 2} \frac{\frac{1}{x^2} - \frac{1}{4}}{x-2}$$

$$\text{Ans.} \quad \lim_{x \rightarrow 2} \frac{(x)^{-2} - (2)^{-2}}{x-2}$$

$$= -2 \times (2)^{-2-1} = -2 \times (2)^{-3} = -2 \times \frac{1}{8} = -\frac{1}{4}$$

$$\textcircled{10} \quad \lim_{x \rightarrow 2} \frac{x-2}{x^4-16}$$

$$\text{Ans.} \quad \lim_{x \rightarrow 2} \frac{x-2}{(x)^4 - (2)^4}$$

$$\lim_{x \rightarrow 2} \frac{x-2}{(x)^4 - (2)^4}$$

$$\lim_{x \rightarrow 2} \frac{x-2}{x^4 - (2)^4} = \frac{1 \times (2)^{4-1}}{(4) \times (2)^{4-1}} = \frac{1}{32}$$

$$\textcircled{11} \quad \lim_{x \rightarrow 0} \frac{(4-x)^4 - 256}{x}$$

$$\text{Ans.} \quad \lim_{x \rightarrow 0} \frac{(4-x)^4 - (4)^4}{x}$$

$$-1 \quad \lim_{x \rightarrow 0} \frac{(4-x)^4 - (4)^4}{4-x-4}$$

$$-1 \quad \lim_{4-x \rightarrow 4} \frac{(4-x)^4 - (4)^4}{(4-x) - 4} = -1 \times 4 \times (4)^{4-1} = -1 \times 4 \times 4^3 = -256$$



⇒ Method of evaluating when  $x \rightarrow \infty$  (Infinity) :-

Formula  $\rightarrow$  Take the highest power of  $x$  common from the numerator and denominator.

Ex  $\Rightarrow$

①  $\lim_{x \rightarrow \infty} \frac{4x+1}{5x-1}$

Ans.  $\lim_{x \rightarrow \infty} \frac{x(4 + \frac{1}{x})}{x(5 - \frac{1}{x})}$

$$\lim_{x \rightarrow \infty} \frac{4 + \frac{1}{x}}{5 - \frac{1}{x}} = \frac{4 + \frac{1}{\infty}}{5 - \frac{1}{\infty}} = \frac{4+0}{5-0} = \frac{4}{5}$$

②  $\lim_{x \rightarrow \infty} \frac{3x^2 + 4x + 5}{x^2 + x}$

Ans.  $\lim_{x \rightarrow \infty} \frac{x^2(3 + \frac{4}{x} + \frac{5}{x^2})}{x^2(1 + \frac{1}{x})}$

$$\lim_{x \rightarrow \infty} \frac{3 + \frac{4}{x} + \frac{5}{x^2}}{1 + \frac{1}{x}} = \frac{3+0+0}{1+0} = \frac{3}{1} = 3$$

③  $\lim_{x \rightarrow \infty} \frac{4x-2}{x^2+x+1}$

Ans.  $\lim_{x \rightarrow \infty} \frac{x(4 - \frac{2}{x})}{x^2(1 + \frac{1}{x} + \frac{1}{x^2})}$

$$\lim_{x \rightarrow \infty} \frac{4 - \frac{2}{x}}{x(1 + \frac{1}{x} + \frac{1}{x^2})} = \frac{4-0}{\infty(1+0+0)} = \frac{4}{\infty} = 0$$

④  $\lim_{x \rightarrow \infty} \frac{3x^2 + x - 1}{2x^2 - 7x + 5}$

Ans.  $\lim_{x \rightarrow \infty} \frac{x^2(3 + \frac{1}{x} - \frac{1}{x^2})}{x^2(2 - \frac{7}{x} + \frac{5}{x^2})}$

$$= \lim_{x \rightarrow \infty} \frac{3 + \frac{1}{x} - \frac{1}{x^2}}{2 - \frac{7}{x} + \frac{5}{x^2}} = \frac{3+0-0}{2-0+0} = \frac{3}{2}$$

$$\textcircled{5} \quad \lim_{x \rightarrow \infty} \frac{1+2+3+\dots+n}{n^2}$$

Ans.  $\lim_{x \rightarrow \infty} \frac{\frac{n(n+1)}{2}}{n^2}$

$$\lim_{x \rightarrow \infty} \frac{n(n+1)}{2n^2}$$

$$\lim_{x \rightarrow \infty} \frac{n^2+n}{2n^2}$$

$$\lim_{x \rightarrow \infty} \frac{\cancel{n^2} \left(1 + \frac{1}{n}\right)}{2\cancel{n^2}} = \frac{1 + \frac{1}{n}}{2} = \frac{1}{2}$$

$$\textcircled{6} \quad \lim_{x \rightarrow \infty} \frac{1^2+2^2+3^2+\dots+n^2}{n^3}$$

Ans.  $\lim_{x \rightarrow \infty} \frac{\frac{n(n+1)(2n+1)}{6}}{n^3}$

$$\lim_{x \rightarrow \infty} \frac{(n^2+n)(2n+1)}{6n^3}$$

$$\lim_{x \rightarrow \infty} \frac{2n^3+n^2+2n^2+n}{6n^3}$$

$$\lim_{x \rightarrow \infty} \frac{2n^3+3n^2+n}{6n^3}$$

$$\lim_{x \rightarrow \infty} \frac{\cancel{n^3} \left(2 + 3\cancel{n} + \frac{1}{n^2}\right)}{6\cancel{n^3}}$$

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{2 + \frac{3}{n} + \frac{1}{n^2}}{6} &= \frac{2+0+0}{6} \\ &= \frac{2}{6} \\ &= \frac{1}{3} \end{aligned}$$



# ⇒ Trigonometric Function :-

$$\textcircled{1} \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

$$\textcircled{2} \lim_{\theta \rightarrow 0} \frac{\theta}{\sin \theta} = 1$$

$$\textcircled{3} \lim_{\theta \rightarrow 0} \frac{\tan \theta}{\theta} = 1$$

$$\textcircled{4} \lim_{\theta \rightarrow 0} \frac{\theta}{\tan \theta} = 1$$

Indeterminate Form :-  $\frac{0}{0}, \frac{\infty}{\infty}$

$$\textcircled{1} 0 \times \infty$$

$$\textcircled{4} 1^\infty$$

$$\textcircled{2} \infty - \infty$$

$$\textcircled{5} \infty^0$$

$$\textcircled{3} 0^0$$

## EX →

$$\textcircled{1} \lim_{x \rightarrow 0} \frac{\sin 2x}{x}$$

Ans.  $\lim_{x \rightarrow 0} \frac{2 \sin 2x}{2x}$

$$2 \lim_{2x \rightarrow 0} \frac{\sin 2x}{2x} = 2 \times 1 = 2$$

$$\textcircled{2} \lim_{x \rightarrow 0} \frac{\sin 5x}{x}$$

Ans.  $\lim_{x \rightarrow 0} \frac{5 \sin 5x}{5x}$

$$5 \lim_{5x \rightarrow 0} \frac{\sin 5x}{5x} = 5 \times 1 = 5$$

$$\textcircled{3} \lim_{x \rightarrow 0} \frac{x}{\sin 10x}$$

Ans.  $\lim_{x \rightarrow 0} \frac{10x}{10 \sin 10x}$

$$\frac{1}{10} \lim_{x \rightarrow 0} \frac{10x}{\sin 10x}$$

$$\frac{1}{10} \lim_{10x \rightarrow 0} \frac{10x}{\sin 10x} = \frac{1}{10} \times 1 = \frac{1}{10}$$

$$\textcircled{4} \lim_{x \rightarrow 0} \frac{\tan 3x}{x}$$

Ans.  $\lim_{x \rightarrow 0} \frac{3 \tan 3x}{3x}$

$$3 \lim_{3x \rightarrow 0} \frac{\tan 3x}{3x} = 3 \times 1 = 3$$

$$\textcircled{5} \lim_{x \rightarrow 0} \frac{\sin x/2}{x}$$

Ans.  $\lim_{x \rightarrow 0} \frac{\frac{1}{2} \sin x/2}{\frac{1}{2} x}$

$$\frac{1}{2} \lim_{x/2 \rightarrow 0} \frac{\sin x/2}{x/2} = \frac{1}{2} \times 1 = \frac{1}{2}$$

$$\textcircled{6} \lim_{x \rightarrow 0} \frac{\tan(4x)}{x}$$

Ans.  $\lim_{x \rightarrow 0} \frac{-4 \tan(-4x)}{-4x}$

$$-4 \lim_{-4x \rightarrow 0} \frac{\tan(-4x)}{-4x} = -4 \times 1 = -4$$

$$\textcircled{7} \lim_{x \rightarrow 0} \frac{\tan 100x}{x}$$

Ans.  $\lim_{x \rightarrow 0} \frac{100 \tan 100x}{100x}$

$$100 \lim_{100x \rightarrow 0} \frac{\tan 100x}{100x} = 100 \times 1 = 100$$

$$\textcircled{8} \lim_{x \rightarrow 0} \frac{\sin 5x}{4x}$$

Ans.  $\lim_{x \rightarrow 0} \frac{\frac{1}{4} \sin 5x}{\frac{1}{4} \times 4x}$

$$\frac{1}{4} \lim_{x \rightarrow 0} \frac{\sin 5x}{x}$$

$$\frac{1}{4} \lim_{x \rightarrow 0} \frac{5 \sin 5x}{5x}$$

$$= \frac{5}{4} \lim_{x \rightarrow 0} \frac{\sin 5x}{5x}$$

$$= \frac{5}{4} \lim_{5x \rightarrow 0} \frac{\sin 5x}{5x}$$

$$= \frac{5}{4} \times 1 = \frac{5}{4}$$

$$⑨ \lim_{x \rightarrow 0} \frac{\sin 3x}{\sin 4x}$$

$$\text{Ans. } \lim_{x \rightarrow 0} \frac{\frac{\sin 3x}{x}}{\frac{\sin 4x}{x}}$$

$$\lim_{x \rightarrow 0} \frac{3 \sin 3x}{3x} \cdot \frac{4 \sin 4x}{4x}$$

$$\frac{3 \lim_{3x \rightarrow 0} \frac{\sin 3x}{3x}}{4 \lim_{4x \rightarrow 0} \frac{\sin 4x}{4x}} = \frac{1 \times 3}{1 \times 4} = \frac{3}{4}$$

$$⑩ \lim_{x \rightarrow 0} \frac{\sin \alpha x}{\tan \beta x}$$

$$\text{Ans. } \lim_{x \rightarrow 0} \frac{\frac{\sin \alpha x}{x}}{\frac{\tan \beta x}{x}}$$

$$\lim_{x \rightarrow 0} \frac{\alpha \frac{\sin \alpha x}{\alpha x}}{\beta \frac{\tan \beta x}{\beta x}}$$

$$\frac{\alpha \lim_{\alpha x \rightarrow 0} \frac{\sin \alpha x}{\alpha x}}{\beta \lim_{\beta x \rightarrow 0} \frac{\tan \beta x}{\beta x}} = \frac{\alpha \times 1}{\beta \times 1} = \frac{\alpha}{\beta}$$

$$⑪ \lim_{x \rightarrow 0} \frac{\tan x/3}{\sin 4x/5}$$

$$\text{Ans. } \lim_{x \rightarrow 0} \frac{\frac{\tan x/3}{x}}{\frac{\sin 4x/5}{x}}$$

$$\lim_{x \rightarrow 0} \frac{\frac{1}{3} \tan x/3}{\frac{4}{5} \sin 4x/5}$$

$$\frac{\frac{1}{3} \lim_{\frac{1}{3}x \rightarrow 0} \frac{\tan x/3}{x/3}}{\frac{4}{5} \lim_{\frac{4}{5}x \rightarrow 0} \frac{\sin 4x/5}{4x/5}} = \frac{\frac{1}{3}}{\frac{4}{5}} = \frac{1}{3} \times \frac{5}{4}$$

$$\frac{\frac{1}{3}}{\frac{4}{5}} = \frac{5}{12}$$

$$⑫ \lim_{x \rightarrow 0} \frac{\sin 4x}{x/2}$$

$$\text{Ans. } \lim_{x \rightarrow 0} \frac{8 \sin 4x}{8x/2}$$

$$8 \lim_{4x \rightarrow 0} \frac{\sin 4x}{4x} = 8 \times 1 = 8$$

$$⑬ \lim_{x \rightarrow 0} \frac{\sin x^\circ}{x}$$

$$\text{Ans. } \lim_{x \rightarrow 0} \frac{\sin \pi x/180}{x}$$

$$\lim_{x \rightarrow 0} \frac{\pi/180 \sin \pi x/180}{\pi/180 x}$$

$$\frac{\pi/180 \lim_{\pi x/180 \rightarrow 0} \frac{\sin \pi x/180}{\pi x/180}}{\pi x/180} = \frac{\pi}{180} \times 1 = \frac{\pi}{180}$$

$$⑭ \lim_{x \rightarrow 0} \frac{\tan 3x}{\sin 4x}$$

$$\text{Ans. } \lim_{x \rightarrow 0} \frac{\frac{\tan 3x}{x}}{\frac{\sin 4x}{x}}$$

$$\lim_{x \rightarrow 0} \frac{3 \tan 3x}{3x} \cdot \frac{4 \sin 4x}{4x}$$

$$\frac{3 \lim_{3x \rightarrow 0} \frac{\tan 3x}{3x}}{4 \lim_{4x \rightarrow 0} \frac{\sin 4x}{4x}} = \frac{3 \times 1}{4 \times 1} = \frac{3}{4}$$

$$180^\circ = \pi$$

$$1^\circ = \frac{\pi}{180}$$

$$x^\circ = \frac{\pi x}{180}$$



$$(15) \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h}$$

Ans. 
$$\lim_{h \rightarrow 0} \frac{2 \cos\left(\frac{x+h+x}{2}\right) \sin\left(\frac{x+h-x}{2}\right)}{h}$$

$$\lim_{h \rightarrow 0} \frac{2 \cos\left(\frac{2x+h}{2}\right) \sin\left(\frac{h}{2}\right)}{h}$$

$$\lim_{h \rightarrow 0} 2 \cos\left(\frac{2x+h}{2}\right) \times \frac{\sin h/2}{2 \times h/2}$$

$$\lim_{h \rightarrow 0} \cos\left(\frac{2x+h}{2}\right) \times \frac{\sin h/2}{h/2}$$

$$\lim_{h \rightarrow 0} \cos\left(\frac{2x+h}{2}\right) \quad \lim_{\frac{h}{2} \rightarrow 0} \frac{\sin h/2}{h/2}$$

$$= \cos x \times 1 = \cos x$$

$$(16) \lim_{h \rightarrow 0} \frac{\cos(x+h) - \cos x}{h}$$

Ans. 
$$\lim_{h \rightarrow 0} \frac{2 \sin\left(\frac{x+h+x}{2}\right) \sin\left(\frac{x-x-h}{2}\right)}{h}$$

$$\lim_{h \rightarrow 0} \frac{2 \sin\left(\frac{2x+h}{2}\right) \sin\left(-\frac{h}{2}\right)}{h}$$

$$- \lim_{h \rightarrow 0} \frac{2 \sin\left(\frac{2x+h}{2}\right) \sin\left(\frac{h}{2}\right)}{h}$$

$$- \lim_{h \rightarrow 0} 2 \sin\left(\frac{2x+h}{2}\right) \frac{\sin h/2}{2 \times h/2}$$

$$- \lim_{h \rightarrow 0} \sin\left(\frac{2x+h}{2}\right) \quad \lim_{h/2 \rightarrow 0} \frac{\sin h/2}{h/2}$$

$$= - \sin x \times 1 = - \sin x$$

$$(17) \lim_{h \rightarrow 0} \frac{\tan(x+h) - \tan x}{h}$$

Ans. 
$$\lim_{h \rightarrow 0} \frac{\frac{\sin(x+h)}{\cos(x+h)} - \frac{\sin x}{\cos x}}{h}$$

$$\lim_{h \rightarrow 0} \frac{\frac{\sin(x+h) \cdot \cos x - \sin x \cdot \cos(x+h)}{\cos x \times \cos(x+h)}}{h}$$

$$\lim_{h \rightarrow 0} \frac{\sin(x+h-x)}{\cos x \times \cos(x+h)}{h}$$

$$\lim_{h \rightarrow 0} \frac{\sin h}{\cos x \times \cos(x+h)} \times \frac{1}{h}$$

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} \times \frac{1}{\cos x \times \cos(x+h)}$$

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} \quad \lim_{h \rightarrow 0} \frac{1}{\cos x \times \cos(x+h)}$$

$$= 1 \times \frac{1}{\cos x \times \cos x} = \frac{1}{\cos^2 x} = \sec^2 x$$

$$(18) \lim_{x \rightarrow 0} \frac{x}{\sin 2x}$$

Ans. 
$$\lim_{x \rightarrow 0} \frac{2x}{2 \sin 2x}$$

$$\frac{1}{2} \lim_{x \rightarrow 0} \frac{2x}{\sin 2x}$$

$$\frac{1}{2} \lim_{2x \rightarrow 0} \frac{2x}{\sin 2x} = \frac{1}{2} \times 1 = \frac{1}{2}$$

$$(19) \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2}$$

Ans. 
$$\lim_{x \rightarrow 0} \frac{2 \sin^2 \frac{x}{2}}{x^2}$$

$$\lim_{x \rightarrow 0} \frac{\sin^2 \frac{x}{2}}{2^{\cancel{2}} \times \frac{x^2}{4}}$$

$$\lim_{x \rightarrow 0} \frac{\sin^2 \frac{x}{2}}{2 \times \frac{x^2}{4}}$$

$$= \frac{1}{2} \lim_{x \rightarrow 0} \left( \frac{\sin \frac{x}{2}}{\frac{x}{2}} \right)^2$$

$$= \frac{1}{2} \lim_{\frac{x}{2} \rightarrow 0} \left( \frac{\sin \frac{x}{2}}{\frac{x}{2}} \right)^2$$

$$= \frac{1}{2} \times (1)^2 = \frac{1}{2}$$



2nd method

$$\rightarrow \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2}$$

$$\lim_{x \rightarrow 0} \frac{(1 - \cos x)(1 + \cos x)}{x^2 (1 + \cos x)}$$

$$\lim_{x \rightarrow 0} \frac{1 - \cos^2 x}{x^2 (1 + \cos x)}$$

$$\lim_{x \rightarrow 0} \frac{\sin^2 x}{x^2} \times \frac{1}{1 + \cos x}$$

$$\lim_{x \rightarrow 0} \left( \frac{\sin x}{x} \right)^2 \lim_{x \rightarrow 0} \frac{1}{1 + \cos x}$$

$$= (1)^2 \times \frac{1}{2} = \frac{1}{2}$$

(20)  $\lim_{x \rightarrow \pi/2} (\sec x - \tan x)$  ( $\infty - \infty$  form)

Ans.  $\lim_{x \rightarrow \pi/2} \left( \frac{1}{\cos x} - \frac{\sin x}{\cos x} \right)$

$$\lim_{x \rightarrow \pi/2} \left( \frac{1 - \sin x}{\cos x} \right)$$

$$\lim_{x \rightarrow \pi/2} \frac{(1 - \sin x)(1 + \sin x)}{\cos x (1 + \sin x)}$$

$$\lim_{x \rightarrow \pi/2} \frac{1 - \sin^2 x}{\cos x (1 + \sin x)}$$

$$\lim_{x \rightarrow \pi/2} \frac{\cos^2 x}{\cos x (1 + \sin x)}$$

$$\lim_{x \rightarrow \pi/2} \frac{\cos x}{1 + \sin x} = \frac{0}{2} = 0$$

2nd method

$$\lim_{x \rightarrow \pi/2} \left( \frac{1}{\cos x} - \frac{\sin x}{\cos x} \right)$$

$$\lim_{x \rightarrow \pi/2} \left( \frac{1 - \sin x}{\cos x} \right)$$

$$\lim_{x \rightarrow \pi/2} \frac{\cos x (1 - \sin x)}{\cos x (\cos x)}$$

$$\lim_{x \rightarrow \pi/2} \frac{\cos x (1 - \sin x)}{\cos^2 x}$$

$$\lim_{x \rightarrow \pi/2} \frac{\cos x (1 - \sin x)}{1 - \sin^2 x}$$

$$\lim_{x \rightarrow \pi/2} \frac{\cos x (1 - \sin x)}{(1 - \sin x)(1 + \sin x)}$$

$$\lim_{x \rightarrow \pi/2} \frac{\cos x}{1 + \sin x} = \frac{0}{2} = 0$$

(21)  $\lim_{x \rightarrow \pi} \frac{1 + \cos x}{\tan^2 x}$  ( $\frac{0}{0}$  form)

Ans.  $\lim_{x \rightarrow \pi} \frac{1 + \cos x}{\frac{\sin^2 x}{\cos^2 x}}$

$$\lim_{x \rightarrow \pi} \frac{(1 + \cos x) \cos^2 x}{\sin^2 x}$$

$$\lim_{x \rightarrow \pi} \frac{(1 + \cos x) \cos^2 x}{1 - \cos^2 x}$$

$$\lim_{x \rightarrow \pi} \frac{(1 + \cos x) \cos^2 x}{(1 + \cos x)(1 - \cos x)}$$

$$\lim_{x \rightarrow \pi} \frac{\cos^2 x}{1 - \cos x} = \frac{1}{2}$$

2nd method

$$\lim_{x \rightarrow \pi} \frac{(1 + \cos x) \cos^2 x}{\sin^2 x}$$

$$\lim_{x \rightarrow \pi} \frac{(1 + \cos x)(1 - \cos x) \cos^2 x}{\sin^2 x (1 - \cos x)}$$

$$\lim_{x \rightarrow \pi} \frac{(1 - \cos^2 x) \cos^2 x}{\sin^2 x (1 - \cos x)}$$

$$\lim_{x \rightarrow \pi} \frac{\sin^2 x \times \cos^2 x}{\sin^2 x (1 - \cos x)}$$

$$\lim_{x \rightarrow \pi} \frac{\cos^2 x}{1 - \cos x} = \frac{1}{2}$$



# ⇒ Exponential / Logarithmic :-

$$\rightarrow \lim_{x \rightarrow 0} (1+x)^{1/x} = e$$

$$\rightarrow \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e$$

$$\rightarrow \lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \ln a$$

$$\rightarrow \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = \ln e = 1$$

$$\rightarrow \log mn = \log m + \log n$$

$$\rightarrow \log \frac{m}{n} = \log m - \log n$$

$$\rightarrow \log m^n = n \log m$$

$$\boxed{\ln = \log e}$$

$$\rightarrow \lim_{x \rightarrow 0} \frac{\log(1+x)}{x} = 1$$

Q1)  $\lim_{x \rightarrow 0} \frac{a^x - 1}{x}$

Ans.  $\ln a$

Q2)  $\lim_{x \rightarrow 0} \frac{2^x - 1}{x}$

Ans.  $\ln 2$

Q3)  $\lim_{x \rightarrow 0} \frac{2^{2x} - 1}{x}$

Ans. 1st method

$$2 \lim_{2x \rightarrow 0} \frac{2^{2x} - 1}{2x} = 2 \ln 2$$

2nd method

$$\lim_{x \rightarrow 0} \frac{(2^2)^x - 1}{x} = \ln(2^2) = 2 \ln 2$$

Q4)  $\lim_{x \rightarrow 0} \frac{3^{2x} - 1}{x}$

Ans.  $\lim_{x \rightarrow 0} \frac{(3^2)^x - 1}{x} = \ln(3^2) = 2 \ln 3$

Q5)  $\lim_{x \rightarrow 0} \frac{4^{10x} - 1}{x}$

Ans.  $10 \lim_{10x \rightarrow 0} \frac{4^{10x} - 1}{10x} = 10 \ln 4$

Q6)  $\lim_{x \rightarrow 0} \frac{2^{5x} - 1}{x}$

Ans.  $\lim_{x \rightarrow 0} \frac{(2^5)^x - 1}{x} = \ln(2^5) = 5 \ln 2$

Q7)  $\lim_{x \rightarrow 0} \frac{2^x - 3^x}{x}$

Ans.  $\lim_{x \rightarrow 0} \frac{2^x - 1 + 1 - 3^x}{x}$

$$\lim_{x \rightarrow 0} \frac{(2^x - 1) - (3^x - 1)}{x}$$

$$\lim_{x \rightarrow 0} \frac{2^x - 1}{x} - \lim_{x \rightarrow 0} \frac{3^x - 1}{x} = \ln 2 - \ln 3$$

Q8)  $\lim_{x \rightarrow 1} \frac{2^{x-1} - 1}{x-1}$

IMP

Ans. Let,  $x-1 = y$   
 $\Rightarrow x = y+1$

$$\lim_{y \rightarrow 0} \frac{2^y - 1}{y}$$

$$\lim_{y \rightarrow 0} \frac{2^y - 1}{y} = \ln 2$$

Q9)  $\lim_{x \rightarrow 0} \frac{a^x - b^x}{x}$

Ans.  $\lim_{x \rightarrow 0} \frac{a^x - 1 + 1 - b^x}{x}$

$$\lim_{x \rightarrow 0} \frac{(a^x - 1) - (b^x - 1)}{x}$$

$$\lim_{x \rightarrow 0} \frac{a^x - 1}{x} - \lim_{x \rightarrow 0} \frac{b^x - 1}{x}$$

$$= \ln a - \ln b$$

$$= \ln \frac{a}{b}$$



⇒ Continuity :- (continuous)

$$\boxed{L.H.L = R.H.L = V.O.L}$$

Q. ① check the continuity  $f(x) = \begin{cases} \frac{\sin x}{\sin 3x} & \text{if, } x \neq 0 \\ \frac{1}{3} & \text{if, } x = 0 \end{cases}$  at  $x = 0$

Ans.

L.H.L

$$\lim_{x \rightarrow 0^-} f(x)$$

$$\lim_{x \rightarrow 0^-} \frac{\sin x}{\sin 3x}$$

$$\therefore (x = 0-h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} \frac{\sin(0-h)}{\sin 3(0-h)}$$

$$\lim_{h \rightarrow 0} \frac{\sin(-h)}{\sin(-3h)}$$

$$\lim_{h \rightarrow 0} \frac{-\sin h}{-\sin 3h}$$

$$\lim_{h \rightarrow 0} \frac{\sin h}{\sin 3h}$$

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} = \frac{1}{3}$$

R.H.L

$$\lim_{x \rightarrow 0^+} f(x)$$

$$\lim_{x \rightarrow 0^+} \frac{\sin x}{\sin 3x}$$

$$\therefore (x = 0+h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} \frac{\sin(0+h)}{\sin 3(0+h)}$$

$$\lim_{h \rightarrow 0} \frac{\sin h}{\sin 3h}$$

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} = \frac{1}{3}$$

$$\lim_{h \rightarrow 0} \frac{\sin h}{h} = \frac{1}{3}$$

V.O.L

$$\text{at } x = 0$$

$$f(x) = \frac{1}{3}$$

$\therefore L.H.L = R.H.L = \text{function exists}$

$L.H.L = R.H.L = V.O.L = \text{continuous.}$

Q. ② check the continuity  $f(x) = \begin{cases} 2x+1 & \text{if, } x < 1 \\ 3x & \text{if, } x = 1 \\ 4x^2-2 & \text{if, } x > 1 \end{cases}$  at  $x = 1$

Ans

L.H.L

$$\lim_{x \rightarrow 1^-} f(x)$$

$$\lim_{x \rightarrow 1^-} (2x+1) \therefore (x = 1-h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} 2(1-h)+1$$

$$\lim_{h \rightarrow 0} 3-2h = 3$$

R.H.L

$$\lim_{x \rightarrow 1^+} f(x)$$

$$\lim_{x \rightarrow 1^+} (4x^2-2) \therefore (x = 1+h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} 4(1+h)^2-2$$

$$= 2$$

V.O.L

$$\text{at } x = 1$$

$$f(x) = 3x = 3 \cdot 1 = 3$$

$L.H.L \neq R.H.L$  (Function

does not exist)

$L.H.L \neq R.H.L \neq V.O.L$

(Discontinuous)

③ check the continuity  $f(x) = \begin{cases} \frac{x^2 - a^2}{x - a} & \text{if } x \neq a \\ a & \text{if } x = a \end{cases}$  at  $x = a$

Ans.

L.H.L

$$\lim_{x \rightarrow a^-} f(x)$$

$$\lim_{x \rightarrow a^-} \frac{x^2 - a^2}{x - a}$$

$$\therefore (x = a - h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} \frac{(a - h)^2 - a^2}{(a - h) - a}$$

$$= 2(a)^{2-1} = 2a$$

R.H.L

$$\lim_{x \rightarrow a^+} f(x)$$

$$\lim_{x \rightarrow a^+} \frac{x^2 - a^2}{x - a}$$

$$\therefore (x = a + h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} \frac{(a + h)^2 - a^2}{(a + h) - a}$$

$$= 2(a)^{2-1} = 2a$$

V.O.L

at  $x = a$

$$f(x) = a$$

$\therefore \text{L.H.L} = \text{R.H.L} = (\text{function exists})$

$\text{L.H.L} = \text{R.H.L} \neq \text{V.O.L} = (\text{discontinuous})$

④ check the continuity  $f(x) = \begin{cases} (1+2x)^{1/x} & \text{if } x \neq 0 \\ e^2 & \text{if } x = 0 \end{cases}$  at  $x = 0$

Ans.

L.H.L

$$\lim_{x \rightarrow 0^-} f(x)$$

$$\lim_{x \rightarrow 0^-} (1+2x)^{1/x}$$

$$\therefore (x = 0 - h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} (1+2(0-h))^{1/(0-h)}$$

$$\lim_{h \rightarrow 0} (1-2h)^{1/-h}$$

$$\lim_{h \rightarrow 0} \left\{ (1+(-2h))^{1/(-2h)} \right\}^2$$

$$= e^2$$

R.H.L

$$\lim_{x \rightarrow 0^+} f(x)$$

$$\lim_{x \rightarrow 0^+} (1+2x)^{1/x}$$

$$\therefore (x = 0 + h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} (1+2(0+h))^{1/h}$$

$$\lim_{h \rightarrow 0} (1+2h)^{1/h}$$

$$\lim_{h \rightarrow 0} \left\{ (1+2h)^{1/(2h)} \right\}^2$$

$$= e^2$$

V.O.L

at  $x = 0$

$$f(x) = e^2$$

$\therefore \text{L.H.L} = \text{R.H.L} = \text{function exist}$

$\text{L.H.L} = \text{R.H.L} = \text{V.O.L} = \text{continuous}$



⑤ check the continuity  $f(x) = \sin \frac{\pi [x]}{2}$  at  $x=0$

Ans.

L.H.L

$$\lim_{x \rightarrow 0^-} f(x)$$

$$\lim_{x \rightarrow 0^-} \sin \frac{\pi [x]}{2}$$

$$\therefore (x = 0-h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} \sin \frac{\pi [0-h]}{2}$$

$$\lim_{h \rightarrow 0} \sin \frac{\pi x(-1)}{2}$$

$$\lim_{h \rightarrow 0} \sin \left(-\frac{\pi}{2}\right)$$

$$-\lim_{h \rightarrow 0} \sin \frac{\pi}{2} = -1$$

R.H.L

$$\lim_{x \rightarrow 0^+} f(x)$$

$$\lim_{x \rightarrow 0^+} \sin \frac{\pi [x]}{2}$$

$$\therefore (x = 0+h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} \sin \frac{\pi [0+h]}{2}$$

$$\lim_{h \rightarrow 0} \sin \frac{\pi x 0}{2}$$

$$\lim_{h \rightarrow 0} \sin \frac{0}{2}$$

$$\lim_{h \rightarrow 0} \sin 0 = 0$$

V.O.L

$$\text{at } x=0$$

$$f(x) = \sin \frac{\pi [x]}{2}$$

$$= \sin \frac{\pi [0]}{2}$$

$$= \sin \frac{0}{2}$$

$$= \sin 0 = 0$$

$\therefore \text{L.H.L} \neq \text{R.H.L}$  (function does not exist)

$\text{L.H.L} \neq \text{R.H.L} = \text{V.O.L}$  (discontinuous)

⑥ If  $f(x) = \begin{cases} ax^2+b & \text{if } x < 1 \\ 1 & \text{if } x = 1 \\ 2ax-b & \text{if } x > 1 \end{cases}$

It is continuous  $x=1$ , find  $a$  and  $b$ .

Ans.

L.H.L

$$\lim_{x \rightarrow 1^-} f(x)$$

$$\lim_{x \rightarrow 1^-} (ax^2+b)$$

$$\therefore (x = 1-h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} \{a(1-h)^2+b\}$$

$$= a+b$$

R.H.L

$$\lim_{x \rightarrow 1^+} f(x)$$

$$\lim_{x \rightarrow 1^+} 2ax-b$$

$$\therefore (x = 1+h, h \rightarrow 0)$$

$$\lim_{h \rightarrow 0} \{2a(1+h)-b\}$$

$$= 2a-b$$

V.O.L

$$\text{at } x=0$$

$$f(x) = 1$$

Since the function is continuous  $\Rightarrow \text{L.H.L} = \text{R.H.L} = \text{V.O.L}$

$$\Rightarrow a+b = 2a-b = 1$$

$$\Rightarrow a+b = 1$$

$$\Rightarrow \frac{2}{3} + b = 1$$

$$\frac{2a+b}{3a+2} = 1$$

$$\Rightarrow \frac{2}{3} + b = 1$$

$$3a+2 \Rightarrow a = \frac{2}{3} \Rightarrow b = 1 - \frac{2}{3} = \frac{1}{3}$$

③ check the continuity

$$f(x) = [3x+11] \quad \text{at } x = -11/3$$

Ans.

L.H.L

$$\lim_{x \rightarrow c^-} f(x)$$

$$x \rightarrow -11/3^-$$

$$\lim_{x \rightarrow -11/3^-} [3x+11]$$

$$(x = (-11/3 - h), h \geq 0)$$

$$\lim_{h \rightarrow 0} [3(-11/3 - h) + 11]$$

$$\lim_{h \rightarrow 0} [-11 - 3h + 11]$$

$$\lim_{h \rightarrow 0} [0 - 3h] = -1$$

R.H.L

$$\lim_{x \rightarrow c^+} f(x)$$

$$x \rightarrow -11/3^+$$

$$\lim_{x \rightarrow -11/3^+} [3x+11]$$

$$\therefore (x = (-11/3 + h), h \geq 0)$$

$$\lim_{h \rightarrow 0} [3(-11/3 + h) + 11]$$

$$\lim_{h \rightarrow 0} [-11 + 3h + 11]$$

$$\lim_{h \rightarrow 0} [0 + 3h] = 0$$

V.O.L

$$\text{at } x = -11/3$$

$$f(x) = [3x+11]$$

$$= [3 \times (-11/3) + 11]$$

$$= [-11 + 11]$$

$$= 0$$

L.H.L  $\neq$  R.H.L (function does not exist)

L.H.L  $\neq$  R.H.L = V.O.L (discontinuous)



## Notes-1

### Topic: DIFFERENTIAL CALCULUS

**DESCRIPTION :** We define the slope of the curve  $y=f(x)$  at the point ,where  $x=a$  to be  $\lim_{h \rightarrow 0} \frac{f(a+h)-f(a)}{h}$ , when it exists this limit is called the derivative of  $f$  at  $x=a$ . now we will look at the derivative as a function derived from  $f$  by considering the limit(slope) at each point of the domain of  $f$ . The derivative of the function " $f$ " with respect to the variable  $x$  is the function " $f'$ " whose value of  $x$  is  $f'(x)=\lim_{h \rightarrow 0} \frac{f(x+h)-f(x)}{h}$ .

A Derivative refers to the instantaneous rate of change of a quantity with respect to the others. That is denoted by  $dy/dx$ , Here  $y=f(x)$ .

Consider the general equation  $f=f(x)$ . Let P&Q be two points of the graph whose abscissas are  $x$  and  $x+h$ . The corresponding ordinates are  $f(x)$  and  $f(x+h)$ . The quantity  $h$ , pictured in below as positive may be either positive or negative. In either case the slope of the secant line P&Q is  $\frac{f(x+h)-f(x)}{x+h-x}$

$$= \frac{f(x+h)-f(x)}{h}$$

Suppose now we keep " $p$ " fixed and let " $Q$ " move along the curve toward " $p$ " (or let  $h$  approach zero). As this happens, the curve may be of such nature that the slope of the secant line varies & approach some fixed value. In that case, the line through  $p$  with slope equal to this limiting value is called the tangent to the curve at  $p$ . Further, the slope of the tangent is said to be the slope of the curve. That is, the slope of the tangent, and also the slope of the curve, at the point  $p(x, y)$  is defined as  $\lim_{h \rightarrow 0} \frac{f(x+h)-f(x)}{h}$  provided the limit exists.

### MODEL QUESTIONS :

**(1)** Find the derivative of the  $X$  at  $x=1$ .

Ans: let  $f(x)=X$  then  $f'(1)=\lim_{h \rightarrow 0} \frac{f(1+h)-f(1)}{h}$

$$= \lim_{h \rightarrow 0} \frac{(1+h)-1}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h}{h}$$

$$= \lim_{h \rightarrow 0} 1$$

$$= 1$$

Thus the derivative at  $x$  at  $x=1$  is 1.

**(2)** Find the derivative of  $x^2$  at  $x=1$

Ans: let  $f(x)=x^2$  then

$$\begin{aligned}f'(1) &= \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} \\&= \lim_{h \rightarrow 0} \frac{[(1+h)^2] - ([1]^2)}{h} \\&= \lim_{h \rightarrow 0} \frac{(1+2.1.h+h^2-1)}{h} \\&= \lim_{h \rightarrow 0} \frac{h(2+h)}{h} \\&= \lim_{h \rightarrow 0} 2 + h = 2\end{aligned}$$

**(3)** Derivative of constant function  $f(x)=c$

$$\begin{aligned}\text{Ans : } f'(c) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h} \\&= \lim_{h \rightarrow 0} \frac{c - c}{h} = 0\end{aligned}$$

### **MOST PROBABLE QUESTIONS:**

- (1) Find the derivative of the  $x^3$  at  $x=1$ .
- (2) Find the derivative of  $x^n$  at  $x=1$
- (1) Find the derivative at  $99x$  at  $x=100$ .
- (2) Find the derivative of  $x^2 - 27$
- (3) Find the derivative of  $\frac{1}{x^2}$
- (4) Find the derivative of  $2x^2 - 2$  at  $x=1$



## Notes-2

### Topic : ALGEBRA OF DERIVATIVE

#### DESCRIPTION :

Now we define the algebra of derivative that is called the laws of derivative .consider two function  $f(x)$  and  $g(x)$  whose derivative in the same domain.

Here we define the algebraic operations of functions like addition ,substraction, multiplication ,scalar multiplication and division

Consider two function  $f(x)$  and  $g(x)$  in the same domain .then their operations

(1) Addition:  $\frac{d}{dx} [f(x) + g(x)] = \frac{d}{dx} f(x) + \frac{d}{dx} g(x)$

(2) Substraction :  $\frac{d}{dx} [f(x) - g(x)] = \frac{d}{dx} f(x) - \frac{d}{dx} g(x)$

(3) Scalar multiplication:  $\frac{d}{dx} [cf(x)] = c \frac{d}{dx} f(x)$

(4) Quotient of two function :  $\frac{d}{dx} \left[ \frac{f(x)}{g(x)} \right] = \frac{f(x) \cdot \frac{d}{dx} [g(x)] - g(x) \cdot \frac{d}{dx} [f(x)]}{[g(x)]^2}$

#### MODEL QUESTIONS :

(1) Find the derivative of function  $f(x) = 2x^2 + 3x + 1$

Ans:  $\frac{d}{dx} (f(x)) = \frac{d}{dx} (2x^2 + 3x + 1)$   
 $= \frac{d}{dx} (2x^2) + \frac{d}{dx} (3x) + \frac{d}{dx} (1)$   
 $= 4x + 3 + 0$   
 $= 4x + 3$

#### MOST PROBABLE QUESTIONS:

(1) Find the derivative of the following

(i)  $8x^3$  (ii)  $5x^2$

(2) find the derivative of the following functions.

(1)  $5x^3 + 2x - 3$

(2)  $3xy$

(3)  $\frac{1}{x}$

(4) find the derivative of the function  $\frac{3xy}{x}$

## Notes -3

### Topic: DERIVATIVE OF STANDARD FUNCTION(Trigonometric function)

#### DESCRIPTION :

Every one has already knows what is trigonometric function in 1<sup>st</sup> semester .it should be kept mind that to find the derivative of trigonometric function ,the angles must be in the radian measure . In case the given angle is measured in degrees, we must first convert it into radian measure by using the formula  $180 \text{ degree} = \pi \text{ radian}$ .

**We** shall now find the derivative of trigonometric function using the definition of the derivative of the function.

(i) Derivative of  $\sin x$ :

Let  $f(x) = \sin x$

Then  $f(x+h) = \sin(x+h)$

Thus  $f(x+h)-f(x) = \sin(x+h)-\sin x$

$$= \frac{f(x+h)-f(x)}{h} = \frac{\sin(x+h)-\sin x}{h} = \frac{2 \cos\left(\frac{2x+h}{2}\right) \sin h/2}{h} = \cos\left(x + \frac{h}{2}\right) \frac{\sin h/2}{h/2}$$

Now taking limit  $h \rightarrow 0$

$$\begin{aligned} &\equiv \lim_{h \rightarrow 0} \frac{f(x+h)-f(x)}{h} = \lim_{h \rightarrow 0} \left[ \cos\left(x + \frac{h}{2}\right) \frac{\sin h/2}{h/2} \right] \\ &= \lim_{h \rightarrow 0} \cos\left(x + \frac{h}{2}\right) \cdot \lim_{h \rightarrow 0} \frac{\sin h/2}{h/2} \\ &= f'(x) = \cos x \quad \left( \text{where } \lim_{h \rightarrow 0} \frac{\sin h/2}{h/2} = 1 \right) \end{aligned}$$

So we get that  $\frac{d}{dx}(\sin x) = \cos x$

(ii) Derivative of  $\cos x$

Let  $f(x) = \cos x$  Then  $f(x+h) = \cos(x+h)$

Since  $f(x+h)-f(x) = \cos(x+h)-\cos x$

$$\text{Or } \frac{f(x+h)-f(x)}{h} = \frac{\cos(x+h)-\cos x}{h} = \frac{-2 \sin\left(x + \frac{h}{2}\right) \sin h/2}{h}$$

$$\text{Since } \lim_{h \rightarrow 0} \frac{f(x+h)-f(x)}{h} = \lim_{h \rightarrow 0} \left[ -\sin\left(x + \frac{h}{2}\right) \frac{\sin h/2}{h/2} \right]$$

$$= -\lim_{h \rightarrow 0} \sin\left(x + \frac{h}{2}\right) \cdot \lim_{h \rightarrow 0} \frac{\sin h/2}{h/2}$$



Then  $f'(\cos x) = -\sin x$

Similarly other functions are define.

$$(iii) \quad \frac{d}{dx}(\tan x) = \sec^2 x$$

$$(iv) \quad \frac{d}{dx}(\operatorname{cosec} x) = -\operatorname{cosec} x \cdot \cot x$$

$$(v) \quad \frac{d}{dx}(\sec x) = \sec x \tan x$$

$$(vi) \quad \frac{d}{dx}(\cot x) = -\operatorname{cosec}^2 x$$

### **MOST PROBABLE QUESTIONS:**

**(1)** find the derivative of following function

$$(i) \quad Y = x^2 \tan x$$

$$\text{Ans: } \frac{d}{dx}(y) = \frac{d}{dx}(x^2 \tan x) = x^2 \cdot \frac{d}{dx}(\tan x) + \tan x \cdot \frac{d}{dx}(x^2) = x^2 \cdot \sec^2 x + \tan x \cdot 2x$$

$$(ii) \quad Y = \sqrt{1 + \sin 2x}$$

$$\text{Ans: } y = \sqrt{1 + \sin 2x} = \sqrt{(\cos x + \sin x)^2} = \cos x + \sin x$$

$$\frac{dy}{dx} = \frac{d}{dx}(\cos x + \sin x) = \frac{d}{dx}(\cos x) + \frac{d}{dx}(\sin x) = -\sin x + \cos x$$

**(2)** find the derivative of the following functions

$$(i) \quad \cot x, \sec x, \operatorname{cosec} x$$

$$(ii) \quad x \sin x$$

$$(iii) \quad 5 \tan x + b \cot x$$

$$(iv) \quad x \cos x + \sin x$$

**(3)** find the derivative of each of the following.

$$(i) \quad \sqrt{\cos x}$$

$$(ii) \quad \frac{1 - \tan x}{1 + \tan x}$$

$$(iii) \quad \frac{\tan x - \cos x}{\sin x \cdot \cos x}$$

$$(iv) \quad \sec^2$$

$$(i) \quad x^{2\frac{1}{x \log_2 e} + \log_2 x} + \sec x \cdot \tan x$$

$$(ii) \quad \frac{-2 \cos x}{(1 + \sin)^2}$$

## Notes -4

### Topic : DERIVATIVE OF EXPONENTIAL FUNCTIONS

#### DESCRIPTION :

The derivative of exponential function are denoted by  $e^x$  or  $a^x$  .

The exponential functions are important point in the derivation form. Which is denoted by

$$\frac{d}{dx}(a^x) = a^x \log_e a \quad \text{and} \quad \frac{d}{dx}(e^x) = e^x.$$

Here we define how to calculate the derivative of exponential function .

(i) Derivative of  $a^x$

Let  $f(x) = a^x$ , then  $f(x+h) = a^{x+h}$

Thus  $f(x+h) - f(x) = a^{x+h} - a^x = a^x(a^h - 1)$

Now  $\frac{f(x+h)-f(x)}{h} = \frac{a^x(a^h-1)}{h}$

Proceeding the limit as h tends to 0, we have

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = a^x \cdot \lim_{h \rightarrow 0} \frac{a^h - 1}{h}$$

$$\therefore \left[ \lim_{h \rightarrow 0} \left( \frac{a^h - 1}{h} \right) = \log_e a \right]$$

$$\therefore f'(x) = a^x \log_e a$$

Similarly other is define.

#### MODEL QUESTIONS :

Find the derivative of the following function with respect to x.

$$(i) \quad \frac{d}{dx}(x^3 + e^x + \cot x) = \frac{d}{dx}(x^3) + \frac{d}{dx}(e^x) + \frac{d}{dx}(\cot x) = 3x^2 + e^x - \operatorname{cosec}^2 x$$

$$(ii) \quad \frac{d}{dx}(\log_e x^3) = \frac{d}{dx}(3 \log_e x) = 3 \cdot \frac{d}{dx}(\log_e x) = \frac{3}{x}$$



**MOST PROBABLE QUESTIONS:**

(1) find the derivative of the following function

(2)  $\frac{3}{\sqrt[3]{x}} - \frac{5}{\cos x} + \log_e x + \frac{6}{\sin x}$

(3)  $3a^x$

(1) find the derivative of each of the function.

(i)  $x^2 - 7$

(ii)  $\frac{1}{2\sqrt{\cos x}} (-\sin x)$

(iii)  $a^x \cdot 2x + \sec x \cdot \tan x$

(iv)  $\frac{a^x(x \ln a - 1) + b^x(1 - x \ln b)}{x^2}$

## Notes -5

### Topic: DERIVATIVE OF LOGARITHMIC FUNCTIONS

#### DESCRIPTION:

As the logarithmic function with base  $a(a>0, a \neq 1)$  and exponential function with the same base form a pair of mutually inverse functions, the derivative of the logarithmic function can also be found by using the inverse function theorem.

First we should know the derivatives for the basic logarithmic functions.

$$\begin{aligned} \text{(i)} \quad \frac{d}{dx} (\ln(x)) &= \frac{1}{x} & \text{(iii)} \quad \frac{d}{dx} (\log_a x) &= \frac{1}{x \log_a x} \\ \text{(ii)} \quad \frac{d}{dx} \log_b x &= \frac{1}{\ln(b) \cdot x} \end{aligned}$$

$$\text{Let } f(x) = \log_a x \therefore f(x+h) = \log_a(x+h)$$

$$\therefore f(x+h) - f(x) = \log_a(x+h) - \log_a x$$

$$= \log_a \left( \frac{x+h}{x} \right) = \log_a \left( 1 + \frac{h}{x} \right)$$

$$\therefore \frac{f(x+h) - f(x)}{h} = \frac{1}{h} \log_a \left( 1 + \frac{h}{x} \right)$$

$$= \frac{1}{x} \cdot \frac{x}{h} \log_a \left( 1 + \frac{h}{x} \right) = \frac{1}{x} \cdot \log_a \left( 1 + \frac{h}{x} \right)^{\frac{x}{h}}$$

$$= \therefore \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \frac{1}{x} \cdot \lim_{h \rightarrow 0} \left( 1 + \frac{h}{x} \right)^{\frac{x}{h}}$$

$$= \frac{1}{x} \cdot \log_a \lim_{h \rightarrow 0} \left( 1 + \frac{h}{x} \right)^{\frac{x}{h}}$$

$$\therefore f'(x) = \frac{1}{x} \log_a e \quad \left[ \text{since } \lim_{h \rightarrow 0} \left( 1 + \frac{h}{x} \right)^{\frac{x}{h}} = e \right]$$

$$\text{Hence we can write as } \log_a e = \frac{1}{\log_e a} \quad [\text{since } \log_a e \cdot \log_e a = 1]$$

$$\text{Thus } \frac{d}{dx} (\log_a x) = \frac{1}{x \log_a a}$$

#### MODEL QUESTIONS : find the derivative of the following problems

$$(1) Y = 2 \ln(3x^2 - 1)$$

Ans: let we put  $u = 3x^2 - 1$  then derivative of  $u$  is given by

$$U' = \frac{du}{dx} = 6x \text{ so the final answer is } \frac{dy}{dx} = 2 \frac{u'}{u} = 2 \times \frac{6x}{3x^2 - 1} = \frac{12x}{3x^2 - 1}$$

$$(2) Y = x(\ln^3 x)$$



Ans: The notation  $y=x(\ln^3 x)$  means  $y=x (\ln x)^3$

This is the product of  $x$  and  $(\ln x)^3$ . so  $\frac{dy}{dx} = x \frac{3(\ln x)^2}{x} + (\ln x)^3(1)$

$$=3(\ln x)^2 + (\ln x)^3$$

$$=(\ln x)^2(3 + \ln x)$$

### **MOST PROBABLE QUESTIONS:**

**find** the derivative of the following functions

(1)  $3 \ln xy + \sin y = x^2$

(2)  $y = (\sin x)^2$  by first taking logarithmic of each side of the equation .

(3)  $y = \ln (\cos x^2)$

find the derivative of the functions

(i)  $y = \log_2 6x$

(ii)  $y = 3 \log_7(x^2 + 1)$

(1)  $Y=\ln \tan \frac{x}{y}$

(2)  $Y=\ln (x + \sqrt{x^2 + a^2})$

(3)  $Y = \ln \left( \frac{1}{\sqrt{1-x^4}} \right)$

## Notes -6

### Topic: DERIVATIVE OF SOME STANDARD FUNCTIONS

#### DESCRIPTION:

We can algebraically find the derivative of all standard function. That is included exponential function, trigonometric function, exponential function, logarithmic function and inverse trigonometric function.

Previously we discuss most of all types of derivative functions. Here we discuss the inverse trigonometry function. Lets discuss some standard formulas .

- (1)  $\frac{d}{dx}(x^n) = nx^{n-1}$
- (2)  $\frac{d}{dx}(a^x) = a^x \log_e a$
- (3)  $\frac{d}{dx}(\log_a x) =$
- (4)  $\frac{d}{dx}(e^x) =$
- (5)  $\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$
- (6)  $\frac{d}{dx}(\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$
- (7)  $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$
- (8)  $\frac{d}{dx}(\cot^{-1} x) = \frac{-1}{1+x^2}$
- (9)  $\frac{d}{dx}(\sec^{-1} x) = \frac{1}{\sqrt{x^2(\sqrt{x^2-1})}}$
- (10)  $\frac{d}{dx}(\csc^{-1} x) = \frac{-1}{\sqrt{x^2(\sqrt{x^2-1})}}$

#### MODEL QUESTIONS: Find the derivative of following functions.

(1)  $y = \sin^{-1} 2x$

Ans:  $\frac{d}{dx}(\sin^{-1} 2x) = \frac{1}{\sqrt{1-(2x)^2}} = \frac{1}{\sqrt{1-4x^2}}$

(2)  $y = \sin^{-1}[x\sqrt{1-x} - \sqrt{x}\sqrt{1-x^2}]$  find  $\frac{dy}{dx}$

Ans: putting  $x = \sin \theta$  and  $\sqrt{x} = \sin \phi$

We get  $y = \sin^{-1}[\sin \theta \cos \phi - \sin \phi \cos \theta]$

$= \sin^{-1}[\sin(\theta - \phi)] = (\theta - \phi) = \sin^{-1} x - \sin^{-1} \sqrt{x}$

$\therefore \frac{dy}{dx} = \frac{d}{dx}(\sin^{-1} x - \sin^{-1} \sqrt{x}) = \frac{d}{dx}(\sin^{-1} x) - \frac{d}{dx}(\sin^{-1} \sqrt{x})$

$= \left[ \frac{1}{\sqrt{1-x^2}} - \frac{1}{2\sqrt{x}\sqrt{1-x}} \right]$



**MOST PROBABLE QUESTIONS:**

**find** the derivative of the followings:

(1)  $Y = \tan^{-1} \sqrt{x}$

(2)  $Y = \cos^{-1}(\cot x)$

(3)  $Y = \cos^{-1}(\tan x)$

differentiate the following functions:

(1)  $Y = \sqrt{\cot^{-1} \sqrt{x}}$

(2) If  $Y = \frac{x \sin^{-1} x}{\sqrt{1-x^2}}$  find  $\frac{dy}{dx}$

(3)  $y = \left( \frac{1-\cos x}{\sin x} \right)$

Find the derivative of the following functions

(1)  $\tan^{-1} \left( \sqrt{\frac{1-\cos x}{1+\cos x}} \right)$

(2)  $\tan^{-1}(\sec x + \tan x)$

(3)  $\cos^{-1} \left( \sqrt{\frac{1+\cos x}{2}} \right)$

## Notes -7

### Topic: DERIVATIVE OF COMPOSITE FUNCTION(CHAIN RULE)

#### DESCRIPTION:

**We** have been differentiating  $y$ , a function of  $x$  with respect to  $x$ . We also comes across since  $u$  when  $y$  is a function of ' $u$ ' and ' $u$ ' is a function of  $x$  that is  $y=f(u)$  and  $u=g(x)$  then  $y=f[g(x)]$  in this case we say  $y$  is a function of a function or  $y$  is a composite function.

We shall now find in method of differentiating composite function.

(chain rule) if  $u$  is a function of  $y$  define by  $y=f(u)$  and  $u$  is a function of  $x$  define by  $u=g(x)$ ,

then  $y$  is a function of  $x$  and  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

Lets more describe about chain rule. Suppose that we have two functions  $f(x)$  and  $g(x)$  and they are both differentiable .

(1) If we define  $F(x) = (f \circ g)(x)$  then the derivative of  $F(x)$  is  $F'(x) = f'(g(x)) g'(x)$

(2) If we have  $y=f(u)$  and  $u=g(x)$  then derivative of  $y$  is  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

Corollary :if  $y=f(u)$  ,  $u=g(v)$  and  $v=h(x)$  then  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx}$  .

By using these formulas we solve the problems.

#### MODEL QUESTIONS : solve these problems by using the chain rule.

(1)  $Y=(2x^3 - 1)^4$  find  $\frac{dy}{dx}$

Ans: let  $u=2x^3 - 1 \therefore y = u^4$  then  $\frac{dy}{du} = 4u^3$  and  $\frac{du}{dx} = 6x^2$

By chain rule ,  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} = 4u^3 \cdot 6x^2 = 24x^2(2x^3 - 1)^3$  .

(2)  $Y=\sqrt{ax^2 + bx + c}$

Ans: let  $u=ax^2 + bx + c$

then  $y=\sqrt{u}$  then  $\frac{dy}{du} = \frac{d}{du}(\sqrt{u}) = \frac{1}{2\sqrt{u}}$

and  $\frac{du}{dx} = \frac{d}{dx}(ax^2 + bx + c) = a \cdot 2x + b \cdot 1 + 0$

$=2ax+b$

By chain rule  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} = \frac{1}{2\sqrt{u}} \cdot (2ax + b) = \frac{2ax+b}{2\sqrt{ax^2+bx+c}}$



**MOST PROBABILITY QUESTIONS:** find  $\frac{dy}{dx}$

(1)  $y = \sqrt{\frac{1-\tan x}{1+\tan x}}$

(2)  $y = \frac{1}{(x^3 + \sin x)^2}$

(3)  $y = \ln(\sqrt{x} + 1)$

(1)  $y = \left(\frac{7x}{x^2+1}\right)^3$  find  $\frac{dy}{dx}$

(2) if  $f(x) = \sin^3 x$ , find  $f'(x)$

(3) find the derivative of  $\sin x^0$

using chain rule to find the derivative of the following.

(1)  $e^{\sin^2 x}$

(2)  $\sqrt{e^{\sqrt{x}}}$

(3)  $\text{Log}(\log x)$

## Notes -8

### TOPIC: DERIVATIVE OF COMPOSITE FUNCTION (CHAIN RULE)

#### DESCRIPTION:

Already we discuss chain rule in previous lecture now we discuss some extra problem., We have been differentiating  $y$ , a function of  $x$  with respect to  $x$ . We also come across since  $u$  when  $y$  is a function of ' $u$ ' and ' $u$ ' is a function of  $x$  that is  $y=f(u)$  and  $u=g(x)$  then  $y=f[g(x)]$  in this case we say  $y$  is a function of a function or  $y$  is a composite function. We shall now find in method of differentiating composite function.

(chain rule) if  $u$  is a function of  $y$  define by  $y=f(u)$  and  $u$  is a function of  $x$  define by  $u=g(x)$ , then  $y$  is a function of  $x$  and  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

Lets more describe about chain rule. Suppose that we have two functions  $f(x)$  and  $g(x)$  and they are both differentiable .

(1) If we define  $F(x) = (f \circ g)(x)$  then the derivative of  $F(x)$  is  $F'(x) = f'(g(x)) g'(x)$

(2) If we have  $y=f(u)$  and  $u=g(x)$  then derivative of  $y$  is  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

Corollary :if  $y=f(u)$  ,  $u=g(v)$  and  $v=h(x)$  then  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx}$  .

By using these formulas we solve the problems .

#### MODEL QUESTIONS :

(1) Differentiate  $\sin^2 x^3$  by using chain rule.

Ans: Let  $y=u^2$  and  $u=\sin x^3$

That is  $y=u^2$ ,  $u=\sin v$  and  $v=x^3$

Applying these chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx}$$

Now  $\frac{dy}{du} = 2u$ ,  $\frac{du}{dv} = \cos v$  and  $\frac{dv}{dx} = 3x^2$

By putting these value to the above equation we get that  $\frac{dy}{dx} = 2u \cdot \cos v \cdot 3x^2 = 6x^2 \sin x^3 \cos x^3$ .



**MOST PROBABLE QUESTIONS:**

using chain rule to find the derivative of each of the following

(1)  $[\tan (3x^2 + 5)]^8$

(2)  $\sqrt{\tan x}$

(3)  $\left(\frac{2 \tan x}{\tan x + \cos x}\right)^2$

(1)  $\log \left(\frac{1-x}{1+x}\right)$

(2)  $\log (x + \sqrt{x^2 + a})$

(3)  $\frac{\log x}{1+x \log x}$

**find the derivative of the following functions by using chain rule.**

(1)  $(3x^2 + 2x + 1)^8$

(2)  $(x^2 + 3)^4$

(3)  $\sin 6x + \cos 7x$

## Notes -9

### TOPIC : DERIVATIVE OF COMPOSITE FUNCTION (CHAIN RULE)

#### DESCRIPTION:

Already we discuss chain rule in previous lecture now we discuss some extra problem. We have been differentiating  $y$ , a function of  $x$  with respect to  $x$ . We also come across since  $u$  when  $y$  is a function of ' $u$ ' and ' $u$ ' is a function of  $x$  that is  $y=f(u)$  and  $u=g(x)$  then  $y=f[g(x)]$  in this case we say  $y$  is a function of a function or  $y$  is a composite function

We shall now find in method of differentiating composite function.

(chain rule) if  $u$  is a function of  $y$  define by  $y=f(u)$  and  $u$  is a function of  $x$  define by  $u=g(x)$  then  $y$  is a function of  $x$  and  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

Lets more describe about chain rule. Suppose that we have two functions  $f(x)$  and  $g(x)$  and they are both differentiable .

(1) If we define  $F(x) = (f \circ g)(x)$  then the derivative of  $F(x)$  is  $F'(x) = f'(g(x)) g'(x)$

(2) If we have  $y=f(u)$  and  $u=g(x)$  then derivative of  $y$  is  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

Corollary :if  $y=f(u)$  ,  $u=g(v)$  and  $v=h(x)$  then  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx}$  .

By using these formulas we solve the problems .

#### MODEL QUESTIONS : differentiate each of the following with respect to $x$ .

(1)  $\log x \cdot e^{\sin x + x^3}$

Ans: let  $y = \log x \cdot e^{\sin x + x^3}$

By using product rule ,

$$\frac{dy}{dx} = \log x \cdot \frac{d}{dx} (e^{\sin x + x^3}) + e^{\sin x + x^3} \cdot \frac{d}{dx} (\log x)$$

$$= \log x \cdot e^{\sin x + x^3} \cdot (\cos x + 3x^2) + e^{\sin x + x^3} \cdot \frac{1}{x}$$

$$= e^{\sin x + x^3} \left[ (\cos x + 3x^2) \log x + \frac{1}{x} \right]$$

(2)  $\log [\log (\log x)]$

Ans: let  $y = \log [\log (\log x)]$

$$\text{Then } \frac{dy}{dx} = \frac{1}{\log (\log x)} \cdot \frac{d}{dx} \log (\log x)$$

$$= \frac{1}{\log (\log x)} \cdot \frac{1}{\log x} \cdot \frac{d}{dx} (\log x)$$

$$= \frac{1}{\log (\log x) \cdot (\log x)} \cdot \frac{1}{x}$$

(3) find the differential coefficient of  $\sin [\cos(\tan x)]$

Ans: let  $y = \sin [\cos (\tan x)]$

$$\frac{dy}{dx} = \cos [\cos (\tan x)] \cdot \frac{d}{dx} [\cos (\tan x)]$$

$$= \cos [\cos (\tan x)] [-\sin (\tan x)] \cdot \frac{d}{dx} (\tan x)$$

$$= -\cos [\cos (\tan x)] [\sin (\tan x)] \cdot \sec^2 x$$

**MOST PROBABLE QUESTIONS:**

using chain rule to find the derivative of each of the following.

(1)  $\log [(\sin x)^{\cos x}]$

(2)  $\sqrt{\cot x}$

(3)  $(3x^4 - 2)$

(4)  $\sqrt{\sin x}$

(5)  $\log (\sin x)$

(6)  $\cos^2 \sqrt{x}$

find the differential coefficient of

(1)  $x^3 \sin^4 x (\log x)^5$

(2)  $\log \{x - 3\sqrt{x^2 - 6x + 1}\}$



## Notes -10

### TOPIC : DIFFERENTIATION

**DESCRIPTION:** Let  $f(x)$  be a real function and  $a$  be any number. Then we define

(i) Right-Hand Derivative:

$\lim_{h \rightarrow 0} \frac{f(a+h)-f(a)}{h}$  if it exists is called the right-hand derivative of  $f(x)$  at  $x=a$  and it

is denoted by  $Rf'(a)$ .

(ii) Left-Hand Derivative:

$\lim_{h \rightarrow 0} \frac{f(a-h)-f(a)}{-h}$  if it exists, is called the left-hand derivative of  $f(x)$  at  $x=a$  and it is denoted by  $Lf'(a)$ .

(Differentiability)

A function  $f(x)$  is said to be differentiable at  $x=a$ , if  $Rf'(a)=Lf'(a)$ . If, however  $Rf'(a) \neq Lf'(a)$ , we say that  $f(x)$  is not differentiable at  $x=a$ .

(Relation between continuity and Differentiability)

Every differentiable function is continuous, but every continuous function is not differentiable.

Proof: let  $f(x)$  be a differentiable function and let  $a$  be any real number in its domain.

$$\text{Then, } \lim_{h \rightarrow 0} \frac{f(a+h)-f(a)}{h} = f'(a)$$

$$\text{Now, } \lim_{h \rightarrow 0} [f(a+h) - f(a)]$$

$$= \lim_{h \rightarrow 0} \left[ \frac{f(a+h)-f(a)}{h} \times h \right]$$

$$= \lim_{h \rightarrow 0} \frac{f(a+h)-f(a)}{h} \times \lim_{h \rightarrow 0} h$$

$$= f'(a) \times 0 = 0$$

$$\text{Thus } \lim_{h \rightarrow 0} [f(a+h) - f(a)] = 0$$

$$\lim_{h \rightarrow 0} f(a+h) = f(a).$$

**MODEL QUESTIONS :** show that the function  $f(x) = x^2$  is differentiable at  $x=1$

**Ans:**  $Rf'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

$$= \lim_{h \rightarrow 0} \frac{(1+h)^2 - (1)^2}{h}$$

$$= \lim_{h \rightarrow 0} \frac{1+h^2+2h-1}{h} = \lim_{h \rightarrow 0} (h+2) = 2$$

$$= \lim_{h \rightarrow 0} \frac{1+h^2-2h-1}{-h} = \lim_{h \rightarrow 0} (-h+2) = 2$$

$$\therefore Rf'(1) = Lf'(1) = 2$$

this show that  $f(x)$  is differentiable at  $x=1$  and  $f'(1)=2$

**MOST PROBABLE QUESTIONS:**

- (1) Show that  $f(x)=[x]$  is not differentiable at  $x=1$
- (2) Show that the constant function is differentiable or not .
- (3) Show that  $f(x)=x$  is differentiable or not at  $x=1$
- (4) Show that the function  $f(x) = \{1+x, \text{ if } x \leq 2 \mid 5-x, \text{ if } x > 2\}$  is not differentiable at  $x=1$ .
- (5) Show that  $f(x) = |x|$  is differentiable or not at  $x=0$ .
- (6) Show that  $f(x) = \log x$  is differentiable or not at  $x=0$ .
- (7) Show that the function  $f(x) = \left\{ x \sin \frac{1}{x}, \text{ when } x \neq 0 \mid 0, \text{ when } x = 0 \right\}$  is continuous but not differentiable at  $x=0$ .
- (8) Show that the function  $f(x) = \left\{ x^2 \cos \frac{1}{x}, \text{ when } x \neq 0 \mid 0, \text{ when } x = 0 \right\}$  is weather continuous or differentiable or both at  $x=0$ .

## Notes-11

**TOPIC :** METHODE OF DIFFERENTIATION (parametric function)

### DESCRIPTION:

Some times both  $x$  and  $y$  may be given as functions of another variable called a parameter.

For example ,any point  $(x, y)$  on the circle  $x^2 + y^2 = r^2$  can be given by  $x = r \cos t, y = r \sin t$ , the variable quantity  $t$  is called parameter. The function consider these variable is called parametric function.

The term parameter is also used to mean a quantity which is invariable for a given curve but changes when we move from the curve of a given type to another. In such case the derivative is given in terms of the variable parameter .In such case the derivative is given in Sterms of the variable parameter

We shall n0w discuss the method of finding  $\frac{dy}{dx}$  when  $x$  and  $y$  are function of  $t$ .

Let  $x = f(t)$  and  $y = g(t)$  corresponding to an increment  $\delta t$  in  $t$ , there are increments  $\delta x$  and  $\delta y$  in  $x$  and  $y$  respectively.

Then  $x + \delta x = f(t + \delta t)$  and  $y + \delta y = g(t + \delta t)$

$$\therefore \delta x = f(t + \delta t) - f(t)$$

And  $\delta y = g(t + \delta t) - g(t)$  from these two equation combine we get

$$\begin{aligned} \frac{\delta y}{\delta x} &= \frac{g(t + \delta t) - g(t)}{f(t + \delta t) - f(t)} = \frac{g(t + \delta t) - g(t)}{f(t + \delta t) - f(t)} \cdot \frac{\delta t}{\delta t} \\ &= \frac{g(t + \delta t) - g(t)}{\delta t} \cdot \frac{\delta t}{f(t + \delta t) - f(t)} \end{aligned}$$

Now as  $\delta t \rightarrow 0, \delta x \rightarrow 0$  and  $\delta y \rightarrow 0$

$$\begin{aligned} \therefore \lim_{\delta t \rightarrow 0} \frac{g(t + \delta t) - g(t)}{\delta t} &\div \lim_{\delta t \rightarrow 0} \frac{f(t + \delta t) - f(t)}{\delta t} \\ \therefore \frac{dy}{dx} &= \frac{dy}{dt} \div \frac{dx}{dt} = \frac{dy/dt}{dx/dt} \end{aligned}$$

### MODEL QUESTIONS :

(1) If  $x = at^2$  and  $y = 2bt$ , find  $\frac{dy}{dx}$

Ans: Here  $x = at^2$  and  $y = 2bt$



$$\frac{dx}{dt} = 2at, \frac{dy}{dt} = 2b \text{ then } \frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = \frac{2b}{2at} = \frac{b}{at}.$$

(2) If  $y = a \cos \theta$  and  $x = a(\theta + \sin \theta)$ , find  $\frac{dy}{dx}$

Ans:  $y = a \cos \theta$  and  $x = a(\theta + \sin \theta)$

$$\frac{dy}{d\theta} = -a \sin \theta, \text{ and } \frac{dx}{d\theta} = a(1 + \cos \theta)$$

$$\frac{dy}{dx} = \frac{dy}{d\theta} \cdot \frac{d\theta}{dx}$$

$$= \frac{-a \sin \theta}{a(1 + \cos \theta)}$$

$$= \frac{-2 \sin \frac{\theta}{2} \cdot \cos \frac{\theta}{2}}{2 \cos^2 \frac{\theta}{2}} = -\tan \frac{\theta}{2}$$

### **MOST PROBABLE QUESTIONS:**

**(1)** Find the derivative of  $x = a \sin^3 t$  and  $y = b \cos^3 t$

**(2)**  $x = \frac{2at}{1+t^2}$  and  $y = \frac{2bt}{1-t^2}$  find the derivative.

**(3)** Find the derivative of the function  $x = \frac{1-t^2}{1+t^2}$  and  $y = \frac{2t}{1+t^2}$

**(4)**  $x = at^2$  and  $y = at^3$

**(5)**  $x = \frac{a(1-t)}{1+t^2}$  and  $y = at(\frac{1-t^2}{1+t^2})$

**(6)**  $x = a\left(\theta + \frac{1}{\theta}\right)$  and  $y = a\left(\theta - \frac{1}{\theta}\right)$

## Notes -12

### TOPIC :DIFFERENTIATION OF PARAMETRIC FUNCTION

#### DESCRIPTION:

Some times both  $x$  and  $y$  may be given as functions of another variable called a parameter.

For example ,any point  $(x, y)$  on the circle  $x^2 + y^2 = r^2$  can be given by  $x = r \cos t, y = r \sin t$ , the variable quantity  $t$  is called parameter. The function consider these variable is called parametric function.

The term parameter is also used to mean a quantity which is invariable for a given curve but changes when we move from the curve of a given type to another. In such case the derivative is given in terms of the variable parameter .In such case the derivative is given in terms of the variable parameter .

We shall now discuss the method of finding  $\frac{dy}{dx}$  when  $x$  and  $y$  are function of  $t$ .

Let  $x = f(t)$  and  $y = g(t)$  corresponding to an increment  $\delta t$  in  $t$ , there are increments  $\delta x$  and  $\delta y$  in  $x$  and  $y$  respectively.

Then  $x + \delta x = f(t + \delta t)$  and  $y + \delta y = g(t + \delta t)$

$$\therefore \delta x = f(t + \delta t) - f(t)$$

And  $\delta y = g(t + \delta t) - g(t)$  from these two equation combine we get

$$\begin{aligned} \frac{\delta y}{\delta x} &= \frac{g(t + \delta t) - g(t)}{f(t + \delta t) - f(t)} = \frac{g(t + \delta t) - g(t)}{f(t + \delta t) - f(t)} \cdot \frac{\delta t}{\delta t} \\ &= \frac{g(t + \delta t) - g(t)}{\delta t} \cdot \frac{\delta t}{f(t + \delta t) - f(t)} \end{aligned}$$

Now as  $\delta t \rightarrow 0, \delta x \rightarrow 0$  and  $\delta y \rightarrow 0$

$$\begin{aligned} \therefore \lim_{\delta t \rightarrow 0} \frac{g(t + \delta t) - g(t)}{\delta t} \div \lim_{\delta t \rightarrow 0} \frac{f(t + \delta t) - f(t)}{\delta t} \\ \therefore \frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt} = \frac{dy/dt}{dx/dt} \end{aligned}$$

#### MODEL QUESTIONS :

(1) find  $\frac{dy}{dx}$ ,  $x = \theta + \sin \theta, y = 1 + \cos \theta$  at  $\theta = \frac{\pi}{4}$

Ans: Here  $x = \theta + \sin \theta$  then  $\frac{dx}{d\theta} = 1 + \cos \theta$

Again  $y = 1 + \cos \theta$  then  $\frac{dy}{d\theta} = -\sin \theta$

$$\frac{dy}{dx} = \frac{dy}{d\theta} \cdot \frac{d\theta}{dx} = \frac{-\sin \theta}{1 + \cos \theta}$$

$$\left[ \frac{dy}{dx} \right] \text{ at } \theta = \frac{\pi}{4} = \frac{-\sin \frac{\pi}{4}}{1 + \cos \frac{\pi}{4}} = \frac{-\frac{1}{\sqrt{2}}}{1 + \frac{1}{\sqrt{2}}} = \frac{-1}{1 + \sqrt{2}}.$$

(2) Find  $\frac{dy}{dx}$ , if  $x = 3 \cot t - 2 \cos^3 t$ ,  $y = 3 \sin t - 2 \sin^3 t$

Ans: Given  $x = 3 \cot t - 2 \cos^3 t$

$$\text{Then } \frac{dx}{dt} = -3 \sin t - 3(\cos^2 t) \cdot (-\sin t)$$

$$= -3 \sin t + 6 \cos^2 t \cdot \sin t = 3 \sin t \cdot \cos 2t$$

Again,  $y = 3 \sin t - 2 \sin^3 t$

$$\text{Then } \frac{dy}{dt} = 3 \cos t - 6 \sin^2 t \cdot \cos t$$

$$= 3 \cos t (1 - 2 \sin^2 t) = 3 \cos t \cdot \cos 2t$$

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = \frac{3 \cos t \cdot \cos 2t}{3 \sin t \cdot \cos 2t}$$

### **MOST PROBABILITY QUESTIONS:**

(1) Find  $\frac{dy}{dx}$ , if  $\sin x = \frac{2t}{1+t^2}$ ,  $\tan y = \frac{2t}{1-t^2}$

(2) find  $\frac{dy}{dx}$  where  $y = x^4 \log x$

(3) What is the derivative of  $x|x|$  at  $x = 2$

(4)  $x = a(\theta + \sin \theta)$  and  $y = a(1 - \cos \theta)$

(5)  $x = \frac{at^2}{1+t^2}$  and  $y = \frac{at^3}{1+t^2}$

(6)  $x = a \sqrt{\frac{t^2-1}{t^2+1}}$  and  $y = at \sqrt{\frac{t^2-1}{t^2+1}}$



## Notes -13

### TOPIC :DIFFERENTIATION OF FUNCTION WITH RESPECT TO FUNCTION

**DESCRIPTION:** IF  $Y = f(X)$  is differentiable , then the derivative of y with respect to x is

$$\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

If  $f$  and  $g$  are differentiable functions of  $x$  and if  $\frac{df}{dg} = \frac{\frac{df}{dx}}{\frac{dg}{dx}} = \frac{f'(x)}{g'(x)}$

The understanding of the differentiation of the function with respect to a function

IF  $Y = f(X)$  is differentiable , then the derivative of y with respect to x is

$$\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

If  $f$  and  $g$  are differentiable functions of  $x$  and if  $\frac{df}{dg} = \frac{\frac{df}{dx}}{\frac{dg}{dx}} = \frac{f'(x)}{g'(x)}$

Which is same as the definition.

Suppose we have two differentiable functions given by  $y = f(x)$  and  $z = g(x)$ . To find the derivative of y with respect to z .we regard x as a parameter and find

$$\frac{dy}{dx} = f'(x), \text{ and } \frac{dz}{dx} = g'(x)$$

i.e.  $\frac{dy}{dz} = \frac{dy}{dx} \cdot \frac{dx}{dz} = \frac{f'(x)}{g'(x)}$

### **MODEL QUESTIONS :**

**(1)** Differentiate  $\tan^{-1} x$  w.r.t  $\tan^{-1} \sqrt{1+x^2}$ .

Ans: Let  $y = \tan^{-1} x$  and  $z = \tan^{-1} \sqrt{1+x^2}$

$$\therefore \frac{dy}{dx} = \frac{1}{1+x^2}, \frac{dz}{dx} = \frac{2x}{2(1+x^2)\sqrt{1+x^2}} = \frac{x}{(2x+x^2)\sqrt{1+x^2}}$$

$$\therefore \frac{dy}{dz} = \frac{dy}{dx} \cdot \frac{dx}{dz}$$

$$= \frac{1}{(1+x^2)} \cdot \frac{(2+x^2)\sqrt{1+x^2}}{x} = \frac{2+x^2}{x\sqrt{1+x^2}}$$

**(2)** Differentiate  $\sin^{-1}(\frac{2x}{1+x^2})$  w.r.t  $\cos^{-1}(\frac{1-x^2}{1+x^2})$

Ans: Set  $x = \tan \theta$  in both the expressions

$$\text{Let } y = \sin^{-1}(\frac{2x}{1+x^2}) \text{ and } z = \cos^{-1}(\frac{1-x^2}{1+x^2})$$

$$Y = \sin^{-1}(\frac{2\tan\theta}{1+\tan^2\theta}) \text{ and } z = \cos^{-1}(\frac{1-\tan^2\theta}{1+\tan^2\theta})$$

$$Y = \sin^{-1}(\sin 2\theta) \text{ and } z = \cos^{-1}(\cos 2\theta)$$

$$Y = 2\theta \text{ and } z = 2\theta$$

$$Y = 2\tan^{-1} x \text{ and } z = 2\tan^{-1} x$$

$$\frac{dy}{dx} = \frac{2}{1+x^2} \text{ and } \frac{dz}{dx} = \frac{2}{1+x^2}$$

$$\frac{dy}{dz} = \frac{dy}{dx} \cdot \frac{dx}{dz} = \frac{2}{(1+x^2)} \cdot \frac{(1+x^2)}{2} = 1.$$

**MOST PROBABLE QUESTIONS:**

- (1) Differentiate  $\sin^2 x$  w.r.t.  $(\ln x)^2$ .
- (2) Differentiate  $e^{\tan x}$  w.r.t.  $\sin x$ .
- (3) Differentiate  $e^{\sin^{-1} x}$  w.r.t.  $e^{-\cos^{-1} x}$

## Notes-14

### TOPIC: DIFFERENTIATION WITH RESPECT TO A FUNCTION

**DESCRIPTION:** IF  $Y = f(X)$  is differentiable, then the derivative of  $y$  with respect to  $x$  is

$$\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

If  $f$  and  $g$  are differentiable functions of  $x$  and if  $\frac{df}{dg} = \frac{\frac{df}{dx}}{\frac{dg}{dx}} = \frac{f'(x)}{g'(x)}$

The understanding of the differentiation of the function with respect to a function

IF  $Y = f(X)$  is differentiable, then the derivative of  $y$  with respect to  $x$  is

$$\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

If  $f$  and  $g$  are differentiable functions of  $x$  and if  $\frac{df}{dg} = \frac{\frac{df}{dx}}{\frac{dg}{dx}} = \frac{f'(x)}{g'(x)}$

Which is same as the definition.

Suppose we have two differentiable functions given by  $y = f(x)$  and  $z = g(x)$ . To find the derivative of  $y$  with respect to  $z$ , we regard  $x$  as a parameter and find

$$\frac{dy}{dx} = f'(x), \text{ and } \frac{dz}{dx} = g'(x)$$

i.e.  $\frac{dy}{dz} = \frac{dy}{dx} \cdot \frac{dx}{dz} = \frac{f'(x)}{g'(x)}$

### MODEL QUESTION:

(1) Differentiate  $\tan^{-1} x$  w.r. to  $\cos^{-1} x$

Ans: let  $y = \tan^{-1} x$  and  $z = \cos^{-1} x$

We have to find  $\frac{dy}{dz}$ . Now

$$\frac{dy}{dx} = \frac{1}{1+x^2} \text{ and } \frac{dz}{dx} = \frac{-1}{\sqrt{1-x^2}}$$

$$\therefore \frac{dy}{dz} = \frac{dy}{dx} \cdot \frac{dx}{dz} = \frac{-\sqrt{1-x^2}}{1+x^2}$$

(2) Differentiate  $e^{\sin x}$  w.r. to  $\cos x$

Ans: let  $y = e^{\sin x}$  and  $z = \cos x$

We have to find  $\frac{dy}{dz} = \frac{dy}{dx} \cdot \frac{dx}{dz} = e^{\sin x} \cdot \cos x \times \frac{-1}{\sin x} = -e^{\sin x} \cdot \cot x$

(3) Differentiate  $\frac{1-\cos x}{1+\cos x}$  w.r. to  $\frac{1-\sin x}{1+\sin x}$

Ans: let  $y = \frac{1-\cos x}{1+\cos x}$  and  $z = \frac{1-\sin x}{1+\sin x}$

$$\text{Now, } \frac{dy}{dx} = \frac{\sin x(1+\cos x) + \sin x(1-\cos x)}{(1+\cos x)^2}$$

$$= \frac{\sin x + \sin x \cos x + \sin x - \sin x \cos x}{(1+\cos x)^2} = \frac{2 \sin x}{(1+\cos x)^2}$$

$$\frac{dz}{dx} = \frac{-\cos(1+\sin x) - \cos x(1-\sin x)}{(1+\sin x)^2} = \frac{-2 \cos x}{(1+\sin x)^2}$$



$$\therefore \frac{dy}{dx} = \frac{dy}{dz} \cdot \frac{dz}{dx}$$

$$\frac{2 \sin x}{(1+\sin x)^2} \cdot \frac{(1+\sin x)^2}{(-2 \cos x)} = -\tan x \frac{(1+\sin x)^2}{(1+\cos x)^2}$$

**MOST PROBABLE QUESTIONS:**

- (1) Differentiate  $\sqrt{x}$  with respect to  $x^2$ .
- (2) Differentiate  $\sin x$  w.r.t  $\cos x$ .
- (3) Differentiate  $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$  w.r.t  $\cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$
- (4) Differentiate  $\left(\frac{\tan^{-1} x}{1+\tan^{-1} x}\right)$  w.r.t  $\tan^{-1} x$ .
- (5) Differentiate  $\tan^{-1}\left(\frac{2x}{1-x^2}\right)$  w.r.t  $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$ .
- (6) Differentiate  $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$  w.r.t  $\tan^{-1}\sqrt{\frac{1-x}{1+x}}$ .
- (7) Differentiate  $\ln(\sin x)$  w.r.t  $\tan x$ .
- (8) Differentiate  $e^{\sin^{-1} x}$  w.r.t  $e^{-\cos^{-1} x}$ .
- (9) Differentiate  $\frac{1-\cos x}{1+\cos x}$  w.r.t  $\frac{1-\sin x}{1+\sin x}$ .

## Notes-15

### TOPIC: DIFFERENTIATION OF IMPLICIT FUNCTION

#### DESCRIPTION:

In mathematics, an implicit equation is a relation of the form  $R(x_1, \dots, x_n) = 0$ , where  $R$  is a function of several variables (often a polynomial). For example, the implicit equation of the unit circle is  $x^2 + y^2 - 1 = 0$ .

Sometimes relationships cannot be represented by an explicit function. For example,  $x^2 + y^2 = 1$ . Implicit differentiation helps us find  $dy/dx$  even for relationships like that. This is done using the chain rule, and viewing  $y$  as an implicit function of  $x$ . For example, according to the chain rule, the derivative of  $y^2$  would be  $2y \cdot (dy/dx)$ .

An implicit function is a function that is defined implicitly by an implicit equation by associating one of the variables (the value) with the others (the arguments). Thus, an implicit function for  $y$  is the context of the unit circle is defined implicitly by  $x^2 + f(x)^2 - 1 = 0$ . The implicit function defines  $f$  as a function of  $x$  only if  $-1 \leq x \leq 1$  and one considers only non-negative (or non-positive) values of the function.

#### MODEL QUESTION:

(1) Find  $\frac{dy}{dx}$ , when  $x^2 + y^2 = 2axy$

Ans: given equation is  $x^2 + y^2 = 2axy$

Differentiating w.r.t.  $x$

$$\frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 2a \frac{d}{dx}(xy)$$

$$\Rightarrow 2x + 2y \frac{dy}{dx} = 2a \left[ x \frac{dy}{dx} + y \right]$$

$$\Rightarrow x + y \frac{dy}{dx} = ax \frac{dy}{dx} + ay \Rightarrow y \frac{dy}{dx} - ax \frac{dy}{dx} = ay - x$$

$$\Rightarrow (y - ax) \frac{dy}{dx} = ay - x \Rightarrow \frac{dy}{dx} = \frac{ay - x}{y - ax}$$

(2) Find  $\frac{dy}{dx}$ ,  $e^y \ln x + \ln y = 0$

Ans: Given equation is  $e^y \ln x + \ln y = 0$

Differentiating w.r.t.  $x$ , we get

$$(e^y) \frac{d}{dx}(\ln x) + \ln x \cdot \frac{d}{dx}(e^y) + x \cdot \frac{d}{dx}(\ln y) + \ln y \cdot \frac{d}{dx}(x) = 0$$

$$\Rightarrow e^y \cdot \frac{1}{x} + \ln x \cdot e^y \cdot \frac{dy}{dx} + x \cdot \frac{1}{y} \frac{dy}{dx} + \ln y = 0$$

$$\Rightarrow \frac{dy}{dx} \left[ \ln x \cdot e^y + \frac{x}{y} \right] = - \left( \frac{e^y}{x} + \ln y \right)$$

$$\Rightarrow \frac{dy}{dx} = - \frac{\left( \frac{e^y + x \ln y}{x} \right)}{\frac{y \ln x \cdot e^y + x}{y}} = - \frac{y(e^y + x \ln y)}{x(y \ln x e^y + x)}$$

**MOST PROBABLE QUESTIOS:**

Find the derivative of y w.r.t x

(1)  $ax^2 + by^2 = 25$

(2)  $\frac{x^2}{9} + \frac{y^2}{16} = 1$

(3)  $e^{xy} + y\sin x = 1$

(4)  $x^3 + y^3 = 3axy$

(5)  $x^y = e^{x-y}$

(6)  $y \tan x - y^2 \cos x + 2x = 0$

(7)  $\tan(x+y) + \tan(x-1) = 0$

(8)  $x^{\frac{1}{2}} y^{-\frac{1}{2}} + x^{\frac{3}{2}} y^{\frac{-3}{2}} = 0$

(9)  $\ln \sqrt{x^2 + y^2} = \tan^{-1}\left(\frac{y}{x}\right)$



## Notes-16

### TOPIC: DIFFERENTIATION OF IMPLICIT FUNCTION

#### DESCRIPTION :

In mathematics, an implicit equation is a relation of the form  $R(x_1, \dots, x_n) = 0$ , where  $R$  is a function of several variables (often a polynomial). For example, the implicit equation of the unit circle is  $x^2 + y^2 - 1 = 0$ .

Sometimes relationships cannot be represented by an explicit function. For example,  $x^2 + y^2 = 1$ . Implicit differentiation helps us find  $dy/dx$  even for relationships like that. This is done using the chain rule, and viewing  $y$  as an implicit function of  $x$ . For example, according to the chain rule, the derivative of  $y^2$  would be  $2y \cdot (dy/dx)$ .

An implicit function is a function that is defined implicitly by an implicit equation by associating one of the variables (the value) with the others (the arguments). Thus, an implicit function for  $y$  is the context of the unit circle is defined implicitly by  $x^2 + f(x)^2 - 1 = 0$ . The implicit function defines  $f$  as a function of  $x$  only if  $-1 \leq x \leq 1$  and one considers only non-negative (or non-positive) values of the function.

#### MODEL QUESTION:

(1) If  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ , find  $\frac{dy}{dx}$ .

Ans: Given that

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0 \dots \dots (1)$$

Differentiating both sides of (i) with respect to  $x$ , we get :

$$2ax + 2h \left\{ x \cdot \frac{dy}{dx} + y \cdot 1 \right\} + 2by \cdot \frac{dy}{dx} + 2g + 2f \cdot \frac{dy}{dx} = 0$$

$$\text{Or } (2ax + 2by + 2g) + (2hx + 2fy + 2f) \cdot \frac{dy}{dx} = 0$$

$$\therefore \frac{dy}{dx} = - \left( \frac{ax + hy + g}{hx + by + f} \right).$$

(1) If  $\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$ , prove that  $\frac{dy}{dx} = \frac{\sqrt{1-y^2}}{\sqrt{1-x^2}}$

Ans: Given that

$$\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y) \dots \dots (1)$$

Putting  $x = \sin \theta$  and  $y = \sin \phi$ , it becomes

$$\cos \theta + \cos \phi = a(\sin \theta - \sin \phi)$$

$$\text{or } \frac{\cos\theta + \cos\phi}{\sin\theta - \sin\phi} = a$$

$$\text{or } \frac{2 \cos\left(\frac{\theta+\phi}{2}\right) \cos\left(\frac{\theta-\phi}{2}\right)}{2 \cos\left(\frac{\theta+\phi}{2}\right) + \sin\left(\frac{\theta-\phi}{2}\right)} = a$$

$$\therefore \cot\left(\frac{\theta - \phi}{2}\right) = a \text{ or } \theta - \phi = 2 \cot^{-1} a$$

$$\text{Thus } \sin^{-1} x - \sin^{-1} y = 2 \cot^{-1} a \dots \dots (ii)$$

$$\frac{1}{\sqrt{1-x^2}} - \frac{1}{\sqrt{1-y^2}} \cdot \frac{dy}{dx} = 0$$

$$\therefore \frac{dy}{dx} = \frac{\sqrt{1-y^2}}{\sqrt{1-x^2}}$$

### **MOST PROBABLE QUESTIONS:**

- (1) Find the derivatives of y w.r.t. x in  $x^2 + y^2 + 5xy = 0$
- (2) Find the derivative of y w.r.t. x in  $\sin^2 x + 2 \cos y + xy = 0$
- (3) Find  $\frac{dy}{dx}$ , where  $\cos(x+y) = y \sin x$
- (4) find  $\frac{dy}{dx}$ , where  $\sin(xy) + \frac{x}{y} = x^2 - y$
- (5) find  $\frac{dy}{dx}$ , where  $(\cos x)^y = (\sin y)^x$
- (6) Find  $\frac{dy}{dx}$ , where  $y \cot x + y^3 \tan x + \sin x = 0$

## Notes-17

### TOPIC: DIFFERENTIATION OF LOGARITHMIC

#### DESCRIPTION :

If we are required to find the differential coefficient of a function whose power is a function of  $x$ , the standard result obtained so far can not be applied directly. In such case we first take the logarithmic of the function and then differentiate. This method is called the logarithmic differentiation.

When the given function is a power of some expression or a product of expression, we take logarithmic on both sides and differentiate the implicit function so obtained.

Now we discuss some rules, by using these things to solve the standard problems.

$$(1) \ y = [f(x)]^{g(x)}$$

$$\begin{aligned} \text{ans: } \log y &= \log [f(x)]^{g(x)} \\ &= g(x) \cdot \log f(x) \end{aligned}$$

$$(2) \ y = [f(x) \cdot g(x)]$$

$$\begin{aligned} \text{ans: } \log y &= \log [f(x) \cdot g(x)] \\ &= \log f(x) + \log g(x) \end{aligned}$$

$$(3) \ y = [f(x)]^{g(x)} + [h(x)]^{v(x)}$$

$$\begin{aligned} \text{ans: let } u &= [f(x)]^{g(x)}, \ v = [h(x)]^{v(x)} \\ y &= u + v \\ \frac{dy}{dx} &= \frac{du}{dx} + \frac{dv}{dx} \end{aligned}$$

#### MODEL QUESTION:

$$(1) \ \text{Find } \frac{dy}{dx}, \text{ when } y = x^x$$

Ans:  $y = x^x$  taking logarithm on both sides, we get

$$\log y = \log x^x = x \log x$$

Differentiating both sides with respect to  $x$ , we get

$$\frac{d}{dx}(\log y) = \frac{d}{dx}(x \log x) = x \cdot \frac{d}{dx}(\log x) + \log x \cdot \frac{d}{dx}(x)$$

$$\Rightarrow \frac{1}{y} \frac{dy}{dx} = x \cdot \frac{1}{x} + \log x = 1 + \log x$$

$$\Rightarrow \frac{dy}{dx} = (1 + \log x)y \Rightarrow \frac{dy}{dx} = x^x(1 + \log x)$$

$$(2) \ \text{Differentiate } (\tan x)^{\sec x}$$

Ans: let  $y = (\tan x)^{\sec x}$

Taking log on both sides

$$\log y = \log(\tan x)^{\sec x}$$



On differentiation ,

$$\frac{1}{y} \cdot \frac{dy}{dx} = \sec x \cdot \frac{\sec^2 x}{\tan x} + \sec x \tan x \cdot \log (\tan x)$$

$$\frac{dy}{dx} = y \left[ \frac{1}{\cos x} \cdot \frac{\cos x}{\sin x} \cdot \sec^2 x + \sec x \cdot \tan x \cdot \log (\tan x) \right]$$

$$= (\tan x)^{\sec x} [\operatorname{cosec} x \cdot \sec^2 x + \sec x \cdot \tan x \log (\tan x)]$$

**MOST PROBABLE QUESTIONS:** Differentiate

(1)  $(\sin x)^x$

(2)  $x^{\sin^{-1} x}$

(3)  $(\sin x)^{\log x}$

(4) Find the derivative of  $(\cos x)^{\ln x} + (\log x)^x$

(5) Find the derivative of  $(\sin x)^{\cos^{-1} x}$

(6) Find the derivative of  $(ax^2 + bx + c)^{\cos x}$

(7) If  $\sqrt{1-x^4} + \sqrt{1-y^4} = k(x^2 - y^2)$  then show that  $\frac{dy}{dx} = \frac{x\sqrt{1-y^4}}{y\sqrt{1-x^4}}$

(8) If  $x^y = e^{x-y}$ , prove that  $\frac{dy}{dx} = \frac{\log x}{(1+\log x)^2}$

(9) Differentiate  $\frac{e^{x^2} \cdot \tan^{-1} x}{\sqrt{1+x^2}}$

## Notes-18

### TOPIC: DIFFERENTIATION OF LOGARITHMIC

#### DESCRIPTION:

If we are required to find the differential coefficient of a function whose power is a function of  $x$ , the standard result obtained so far can not be applied directly. In such case we first take the logarithmic of the function and then differentiate. This method is called the logarithmic differentiation.

When the given function is a power of some expression or a product of expression, we take logarithmic on both sides and differentiate the implicit function so obtained. Here we discuss more problems related to logarithmic function.

Now we discuss some rules, by using these things to solve the standard problems.

$$(4) \ y = [f(x)]^{g(x)}$$

$$\begin{aligned} \text{ans: } \log y &= \log [f(x)]^{g(x)} \\ &= g(x) \cdot \log f(x) \end{aligned}$$

$$(5) \ y = [f(x) \cdot g(x)]$$

$$\begin{aligned} \text{ans: } \log y &= \log [f(x) \cdot g(x)] \\ &= \log f(x) + \log g(x) \end{aligned}$$

$$(6) \ y = [f(x)]^{g(x)} + [h(x)]^{v(x)}$$

$$\text{ans: let } u = [f(x)]^{g(x)}, v = [h(x)]^{v(x)}$$

$$y = u + v$$

$$\frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$$

#### MODEL QUESTION:

$$(1) \text{ If } y = x^{x^x}, \text{ find } \frac{dy}{dx}$$

$$\text{Ans: let } y = x^{x^x} \text{ then } \log y = x^x (\log x)$$

On differentiating both sides with respect to  $x$ , we get :

$$\frac{1}{y} \cdot \frac{dy}{dx} = x^x \cdot \frac{1}{x} + (\log x) \cdot \frac{d}{dx} (x^x)$$

$$\therefore \frac{dy}{dx} = y [x^{x-1} + (\log x) \cdot x^x (1 + \log x)]$$

$$[\because u = x^x \Rightarrow \log u = x \log x]$$

$$\Rightarrow \frac{1}{u} \cdot \frac{du}{dx} = (x \cdot \frac{1}{x} + (\log x) \cdot 1) \Rightarrow \frac{du}{dx} = u(1 + \log x) = x^x (1 + \log x)$$

$$\text{hence, } \frac{dy}{dx} = x^{x^x} [x^{x-1} + (\log x) x^x (1 + \log x)]$$

(2) Differentiate  $\tan x \tan 2x \tan 3x \tan 4x$

Ans:

$$\text{Let } y = \tan x \tan 2x \tan 3x \tan 4x$$

$$\text{Then, } \log y = \log (\tan x) + \log (\tan 2x) + \log (\tan 3x) + \log (\tan 4x)$$

$$\frac{1}{y} \cdot \frac{dy}{dx} = \left\{ \frac{\sec^2 x}{\tan x} + \frac{2 \sec^2 2x}{\tan 2x} + \frac{3 \sec^2 3x}{\tan 3x} + \frac{4 \sec^2 4x}{\tan 4x} \right\}$$

$$\therefore \frac{dy}{dx} = y \left[ \frac{1}{\sin x \cos x} + \frac{2}{\sin 2x \cos 2x} + \frac{3}{\sin 3x \cos 3x} + \frac{4}{\sin 4x \cos 4x} \right]$$

$$= y \left[ \frac{2}{\sin 2x} + \frac{4}{\sin 4x} + \frac{6}{\sin 6x} + \frac{8}{\sin 8x} \right]$$

$$= [2 \tan x \tan 2x \tan 3x \tan 4x] \times [\operatorname{cosec} 2x + 2 \operatorname{cosec} 4x + 3 \operatorname{cosec} 6x + 4 \operatorname{cosec} 8x]$$

**MOST PROBABLE QUESTIONS:** Differentiate

(1)  $x^x$

(2)  $(\ln x)^x$

(3)  $x^{x^2}$

(4)  $x^{\ln x}$

(5)  $x^{\tan x} + \cos x^{\sin x}$

(6)  $\sqrt{x(x+1)(x+2)}$

(7)  $x^{\sin x} + \tan x^x$

(8) if  $y = \left( \sqrt{x}^{\sqrt{x}^{\sqrt{x}}} \right)$ , prove that  $\left( \frac{dy}{dx} \right) = \frac{y^2}{(2-y \log x)}$

(9) if  $y = x^{x^{x^{\dots \infty}}}$ , prove that  $\frac{dy}{dx} = \frac{y^2}{x(1-y \log x)}$

(10) if  $y = \sqrt{\sin x + \sqrt{\sin x + \dots \infty}}$ , prove that  $\frac{dy}{dx} = \frac{\cos x}{(2y-1)}$



## Notes-19

**TOPIC:**APPLICATION OF DERIVATIVE(Successive differentiation ( second order)

### DESCRIPTION:

Let  $f(x)$  be a function, differentiable on an open interval  $(a,b)$  . then we know  $f'(x)$  exist at each  $x \in (a,b)$ . The correspondence  $x \rightarrow f'(x), x \in (a,b)$  is a function in its own right . This new function is denoted by  $f'(x)$  and is called derivative of  $f(x)$ .

If the derivative  $f'(x)$  of a function  $f(x)$  is itself differentiable then the derivative of  $f'(x)$  is called the second order derivative of  $f(x)$  and is denoted by  $f''(x)$ .

If  $y = f(x)$  then  $\frac{dy}{dx}$ , the derivative of  $y$  w.r.t  $x$  , is itself , in general , a function of  $x$  and can be differentiable again. To fix up the idea , we shall call  $\frac{dy}{dx}$  as the first order derivative of  $y$  with respect to  $x$  and the derivative of  $\frac{dy}{dx}$  w.r.t  $x$  as second order derivative of  $y$  w.r.t  $x$  and will denoted by  $\frac{d^2y}{dx^2}$  similarly the higher order derivative is denoted by  $\frac{d^ny}{dx^n}$ .

If  $y = f(x)$  then the order alternative notation for  $\frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots, \frac{d^ny}{dx^n}$  . that is also denoted by  $f'(x), f''(x), \dots, f^n(x)$ . Which is denoted by  $f_1, f_2, \dots, f_n$ .

### MODEL QUESTION:

(1) If  $x = at^2, y = 2at$  find  $\frac{d^2y}{dx^2}$

Ans: Given  $x = at^2, y = 2at$

$$\begin{aligned} \frac{dx}{dt} &= 2at, \frac{dy}{dt} = 2a, \therefore \frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = 2a \cdot \frac{1}{2at} = \frac{1}{t} \\ \therefore \frac{d^2y}{dx^2} &= \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d}{dx} \left( \frac{1}{t} \right) = \frac{d}{dt} \left( \frac{1}{t} \right) \cdot \frac{dt}{dx} = -\frac{1}{t^2} \cdot \frac{1}{2at} = -\frac{1}{2at^3} \end{aligned}$$

(2) If  $y = \tan^{-1} x$  , prove that  $(1+x^2)y_2 + 2xy_1 = 0$

Ans: Given  $y = \tan^{-1} x, y_1 = \frac{1}{1+x^2}$

$$\Rightarrow (1+x^2)y_1 = 1$$

Again differentiating w.r.t.  $x$

$$\Rightarrow (1+x^2) \cdot \frac{d}{dx}(y_1) + y_1 \frac{d}{dx}(1+x^2) = 0$$

$$\Rightarrow (1+x^2)y_2 + y_1 \cdot 2x = 0$$

$$\Rightarrow (1+x^2)y_2 + 2xy_1 = 0$$

(3) If  $y = A \cos nx + B \sin nx$  then show that  $\frac{d^2y}{dx^2} + n^2y = 0$

Ans: Given  $A \cos nx + B \sin nx$

$$\frac{dy}{dx} = -A \sin nx \cdot n + B \cos nx \cdot n$$

$$\frac{d^2y}{dx^2} = -An \cdot \cos nx \cdot n - Bn \cdot \sin nx \cdot n = -n^2(A \cos nx + B \sin nx)$$

$$\frac{d^2y}{dx^2} = -ny^2 \Rightarrow \frac{d^2y}{dx^2} + n^2y = 0$$

### **MOST PROBABLE QUESTIONS:**

- (1) What is the slope of the curve  $y = \sin x$  at  $x = \frac{\pi}{6}$
- (2) If  $y = x \sin x$ , what is  $y_1$ , at  $x = 0$
- (3) Find the 2<sup>nd</sup> derivative of the function  $\cos 2x$ .
- (4) If  $x = 2 \cos t - \cos 2t$ ,  $y = 2 \sin t - \sin 2t$ , find  $\frac{d^2y}{dx^2}$ .
- (5) If  $x = a(\theta - \sin \theta)$ ,  $y = a(1 + \cos \theta)$ , find  $\frac{dy}{dx}, \frac{d^2y}{dx^2}$
- (6) If  $y = a \sin 2x + b \cos 2x$ , show that  $\frac{d^2y}{dx^2} + 4y = 0$
- (7) If  $y = \sin(\sin x)$  prove that  $\frac{d^2y}{dx^2} + \tan x \frac{dy}{dx} + y \cos^2 x = 0$
- (8) If  $Y = A \cos nx + b \sin nx$  then show that  $\frac{d^2y}{dx^2} + n^2y = 0$ .
- (9) If  $y = (\sin^{-1} x)^2$ , show that  $(1-x^2)y_2 - xy_1 - 2 = 0$ .

## Notes -20

**TOPIC:**PARTIAL DIFFERENTIAL EQUATION(Function of two variables second order)

**DESCRIPTION:** So far we have studied about derivatives of function of a single variable i.e  $y=f(x)$

$\frac{dy}{dx} = \lim_{\delta x \rightarrow 0} \frac{f(x+\delta x) - f(x)}{\delta x} = f'(x)$ , in that case if a dependent variable is a function of single independent variable .in order to find the derivative of a function of two variables the following procedure is adopted

If a dependent variable is a function of two or more independent variables, in that case partial derivatives exists i.e  $z=f(x,y)$ .The function is differentiated with respect to one of the independent variables while other is treated as constant.

Consider a function of two independent variables  $x$  and  $y$ . Let  $z=f(x,y)$  If the variable  $x$  under goes a change  $\delta x$ .let the variable  $y$  remains constant, then  $z$  undergoes a change  $\delta z$

$$\delta z = f(x + \delta x, y) - f(x, y)$$

We say that  $z$  possess partial derivative w.r.t.  $x$  and denoted by

$$\frac{\partial z}{\partial x} = \lim_{\delta x \rightarrow 0} \frac{f(x + \delta x, y) - f(x, y)}{\delta x} = \frac{\partial f}{\partial x} = f_x$$

Similarly  $z$  possesses partial derivative w.r.t.  $y$  and denoted by

$$\frac{\partial z}{\partial y} = \lim_{\delta y \rightarrow 0} \frac{f(x, y + \delta y) - f(x, y)}{\delta y} = \frac{\partial f}{\partial y} = f_y$$

Let  $z=f(x,y)$  be a function of two variables. Then

$\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$  are themselves functions of two variables  $x$  and  $y$ ,  $\frac{\partial z}{\partial x} = p$ ,  $\frac{\partial z}{\partial y} = q$

$$= \frac{\partial^2 f}{\partial y^2} = f_{yy} = t \frac{\partial^2 z}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial z}{\partial x} \right) = \frac{\partial^2 f}{\partial x^2} = f_{xx} = r \frac{\partial^2 z}{\partial y^2} = \frac{\partial}{\partial y} \left( \frac{\partial z}{\partial y} \right)$$

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial}{\partial x} \left( \frac{\partial z}{\partial y} \right) = f_{xy} = s, \frac{\partial^2 z}{\partial y \partial x} = \frac{\partial}{\partial y} \left( \frac{\partial z}{\partial x} \right) = f_{yx} = s$$

In general  $\left( \frac{\partial^2 z}{\partial xy} \right) \neq \frac{\partial^2 z}{\partial y \partial x}$ .

### **MODEL QUESTION:**

(1) If  $z = x^2y + xy^2$ , find  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$

Ans:  $z = x^2y + xy^2$

$$\frac{\partial z}{\partial x} = \frac{\partial}{\partial x} (x^2y + xy^2) = \frac{\partial}{\partial x} (x^2y) + \frac{\partial}{\partial x} (xy^2) = 2xy + y^2$$



$$\frac{\partial z}{\partial y} = \frac{\partial}{\partial y}(x^2y + xy^2) = \frac{\partial}{\partial y}(x^2y) + \frac{\partial}{\partial y}(xy^2) = x^2 + 2xy$$

(2) If  $z = \sin\left(\frac{x}{y}\right)$  find  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$

Ans: Given  $z = \sin\left(\frac{x}{y}\right)$

$$\frac{\partial z}{\partial x} = \frac{\partial}{\partial x}\left\{\sin\left(\frac{x}{y}\right)\right\} = \cos\left(\frac{x}{y}\right) \frac{\partial}{\partial x}\left(\frac{x}{y}\right) = \frac{1}{y} \cos\left(\frac{x}{y}\right)$$

$$\frac{\partial z}{\partial y} = \frac{\partial}{\partial y}\left\{\sin\left(\frac{x}{y}\right)\right\} = \cos\left(\frac{x}{y}\right) \cdot \frac{\partial}{\partial y}\left(\frac{x}{y}\right) = \cos\left(\frac{x}{y}\right) \cdot \left(-\frac{x}{y^2}\right) = -\frac{x}{y^2} \cos\left(\frac{x}{y}\right)$$

(3) If  $f(x, y) = \frac{2x-3y}{x^2+y^2}$ , find  $f_x(1,2)$  and  $f_y(1,2)$ .

Ans:  $f(x, y) = \frac{2x-3y}{x^2+y^2}$ . differentiate f w.r.t x, treating y as constant, we have

$$f_x(x, y) = \frac{(x^2+y^2) \cdot 2 - (2x-3y) \cdot 2x}{(x^2+y^2)^2} = \frac{6xy - 2x^2 + 2y^2}{(x^2+y^2)^2}$$

$$\therefore f_x(1,2) = \frac{6 \cdot 1 \cdot 2 - 2 \cdot 1^2 + 2 \cdot 2^2}{(1^2 + 2^2)^2} = \frac{18}{25}$$

Differentiating f w.r.t y, treating x as constant, we have

$$f_y(x, y) = \frac{(x^2+y^2) \cdot (-3) - (2x-3y) \cdot 2y}{(x^2+y^2)^2} = \frac{3y^2 - 3x^2 - 4xy}{(x^2+y^2)^2}$$

$$\therefore f_y(1,2) = \frac{3 \cdot 2^2 - 3 \cdot 1^2 - 4 \cdot 1 \cdot 2}{(1^2 + 2^2)^2} = \frac{1}{25}$$

### **MOST PROBABLE QUESTIONS:**

find  $f_x, f_y$  where  $f(x, y)$  is given by

(1)  $\frac{x^2y + xy^2}{x+y}$

(2)  $x^y + y^x$

(3)  $\sin^{-1}\left(\frac{x}{y}\right)$

(4) If  $f(x, y, z) = e^{xyz}$  then find  $xf_x + yf_y + zf_z$

(5) If  $u = (x^2 + y^2 + z^2)^{-\frac{1}{2}}$  show that  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0$

(6) If  $z = f\left(\frac{y}{x}\right)$ , show that  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 0$

(7) Find the degree of the homogeneous function  $f(x, y) = x^4 + x^3y - y^4$ , by two different methods.

(8) If  $z = \tan^{-1}\left(\frac{x^3+y^3}{x+y}\right)$  show that  $x \frac{\partial z}{\partial x} + \frac{\partial z}{\partial y} = \sin 2z$ .

(9) If  $z = \sin^{-1}\left(\frac{xy}{x+y}\right)$  show that  $x \frac{\partial z}{\partial x} + \frac{\partial z}{\partial y} = \tan x$

## Notes -21

**TOPIC:** PROBLEM BASED ON ABOVE

### MOST PROBABLE QUESTIONS:

- (1) If  $f(x, y) = e^{xy}$  what is  $y \frac{\partial f}{\partial y} - x \frac{\partial f}{\partial x}$
- (2) If  $u = \sin x \cos y$ , what is  $\frac{\partial u}{\partial y}$
- (3) What is the slope of the curve  $y = \sin x$  at  $x = \frac{\pi}{6}$
- (4) What is the slope of the tangent to the curve  $y = \sin x$  at  $x = \frac{\pi}{3}$
- (5) Find the slope of the curve  $y = \frac{5}{3}x^2$  at  $x = 2$
- (6) Find the derivative of the following functions
  - (i)  $(x^3 + e^x + 3^x + \cot x)$
  - (ii)  $(9x^2 + \frac{3}{x} + 5 \sin x)$
  - (iii)  $(x^2 + \frac{4}{x^2} - \frac{2}{3} \tan x + 7 \log_e x + 6e)$
  - (iv)  $\log_e x^3$
- (7) Given  $y = (2x^3 - 4)^5$ , find  $\frac{dy}{dx}$
- (8) Find the derivative of function
  - (i)  $y = e^{\sin x}$
  - (ii)  $y = \log (\sin x)$
- (9) Find the derivative of  $\sin^{-1} 5x$
- (10) Find the derivative of  $\tan^{-1} \sqrt{x}$

### FIVE MARK QUESTION:

- (1) Differentiate  $\frac{(1-x)^{\frac{1}{2}}(2-x^2)^{\frac{2}{3}}}{(3-x^3)^{\frac{3}{4}}(4-x^4)^{\frac{4}{5}}}$
- (2) Differentiate  $e^{\sin^{-1} x}$  w.r. to  $e^{-\cos^{-1} x}$
- (3) Differentiate  $\ln (\sin x)$  w.r. to  $\tan x$
- (4) If  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ , find  $\frac{dy}{dx}$
- (5) If  $\sin \sin y = x \sin (a + y)$ , prove that  $\frac{dy}{dx} = \frac{\sin^2(a+y)}{\sin a}$
- (6) Find the tangent line to  $f(x) = 4\sqrt{2x} - 6e^{2-x}$  at  $x=2$
- (7) Find the derivative of  $f(x) = \frac{1+e^{-2x}}{x+\tan(12x)}$  using chain rule.
- (8) Find the derivative  $h(u) = \tan(4 + 10u)$  by using chain rule.
- (9) Find the derivative of  $u(t) = \tan^{-1}(3t - 1)$  by using chain rule.
- (10) Find the first derivative of  $f(x) = (\sqrt{x} + 2x)(4x^2 - 1)$

